

تم تحميل هذا الملف من موقع المناهج الإماراتية



## أسئلة مراجعة نهاية الفصل وفق الهيكل الوزاري الخطة 102A-M بعد التعديل

موقع المناهج ← المناهج الإماراتية ← الصف الثاني عشر المتقدم ← فيزياء ← الفصل الأول ← ملفات المدرس ← الملف

تاريخ إضافة الملف على موقع المناهج: 2024-11-08 21:51:51

ملفات اكتب للمعلم اكتب للطالب | اختبارات الكترونية | اختبارات | حلول | عروض بوربوينت | أوراق عمل  
منهج انجليزي | ملخصات وتقارير | مذكرات وبنوك | الامتحان النهائي للمدرس

المزيد من مادة  
فيزياء:

إعداد: عبد الرحمن عصام

## التواصل الاجتماعي بحسب الصف الثاني عشر المتقدم



الرياضيات



اللغة الانجليزية



اللغة العربية



التربية الاسلامية



المواد على تلغرام

صفحة المناهج  
الإماراتية على  
فيسبوك

## المزيد من الملفات بحسب الصف الثاني عشر المتقدم والمادة فيزياء في الفصل الأول

مراجعة القسم الالكتروني الاختياري وفق الهيكل الوزاري

1

حل أسئلة الامتحان النهائي القسم الالكتروني

2

ملزمة شرح وتدرجات الوحدة الرابعة Capacitors المكثفات

3

أسئلة مراجعة نهاية الفصل وفق الهيكل الوزاري الخطة 102A-M

4

المزيد من الملفات بحسب الصف الثاني عشر المتقدم والمادة فيزياء في الفصل الأول

أسئلة الوحدة الثانية Electric Field The وفق الهيكل الوزاري الخطة C-102

5

# استبيان



EOT I2ADV — M.I02-A

مع خالص الدعاء بالتوفيق والنجاح

أ/ عبد الرحمن عصام

0509886279



Term 1

12 Final revision

Grade 12 Advanced — M.102-A

EOT.A 2025

|   |  |
|---|--|
| Academic Year                                   | 2024/2025                                      |
| Term  | 1  |
| Subject   | Physics M 102 A                                |
| Grade   | 12   |
| Stream  | Advanced                                       |
| Number of Electronic Questions (Swift Assess)   | 15   |
| Mark per Question                               | 4  |
| Number of Free Responses Questions (Paper Part) | 4  |
| Mark per Question                               | 10   |
| Type of All Questions                           | Electronic Questions & Free Response Questions |
| Maximum Overall Grade                           | 100  |
| Exam Duration                                   | 150 minutes                                    |
| Mode of Implementation                          | Swift Assess & Paper Part                      |
| Calculator                                      | Allowed  |

| Question*   | Learning Outcomes Or KPIs**  | Reference(s) in the Student Book (Arabic / English Version)                     |                                  |
|---|--|---|----------------------------------|
|   |  | Example/Exercise  | Page                             |
| 1   | Apply Coulomb's law to relate the magnitude of the electrostatic force, the charge magnitudes of the pair of interacting particles, and the separation between them.   | EXAMPLE 1.2<br>EXAMPLE 1.3<br>SOLVED PROBLEM 1.1<br>EXERCISES 1.83 & 1.84 p: 25 | 10, 11,<br>12, 13 & 14           |
| 2<br>3  | Solve problems involving general charge distribution and the electric field<br>Develop a tool, sketches, descriptive text or presentation to show the morphology of electric field lines of a single or multiple charge system with positive and/or negative charges<br>Define the electric field vector at a point in space as the quotient of the electrostatic force vector acting on a positive test charge at this point divided by the test charge   | As mentioned in the book SOLVED PROBLEM 2.2                                     | 28, 29 & 30<br>34, 35 & 36       |
| 4<br>5  | Find for a uniform distribution of charge, the linear charge density $\lambda$ for charging along a line, the surface charge density $\sigma$ for charge on a surface, and the volume<br>Define the electric field vector at a point in space as the quotient of the electrostatic force vector acting on a positive test charge at this point divided by the test charge  | As mentioned in the book SOLVED PROBLEM 2.2                                     | 28, 29 & 30<br>34, 35 & 36<br>53 |
| 6<br>7  | Apply the relationship between the electric field $E$ and the electric force $F$ and the charge $q$  | As mentioned in the book<br>MULTIPLE-CHOICE QUESTIONS 2.5                       | 28, 29 & 30<br>34, 35 & 36<br>53 |
| 8   | Apply the relationship between the electric field $E$ and the electric force $F$ and the charge $q$  | As mentioned in the book SOLVED PROBLEM 2.2                                     | 37, 38 & 39                      |
| 9<br>10   | Solve problems on electric flux Define the electric flux through a surface as the dot product between the electric field vector and the area vector at each point of that surface and expresses that in an equation<br>Prove that the electric flux through a closed surface is given by the net charge inside the surface divided by the permittivity of the medium, and write the Gauss's law in its integral form   | FIGURE 2.22<br>FIGURE 2.23  | 42, 43 & 44                      |
| 11  | Apply the relationship between the charge density and the electric field magnitude $E$ and also specify the direction of the field for points near a flat, thin, infinite or large, nonconducting/conducting surface with a uniform charge density   | As mentioned in the book  | 47, 48 & 49                      |
| 12  | Solve problems involving electric potential energy   | As mentioned in the book FIGURE 3.2   | 60 & 61                          |
| 13  | Develop a method such as schematic representations to compare the equipotential surfaces due to a point charge, two identical charges, and two different charges   | FIGURE 3.17<br>FIGURE 3.18<br>FIGURE 3.19                                       | 67, 68 & 69                      |
| 14  | Relate the component of the electric field along a certain direction $E_x$ to the change in the electric potential along that direction ( $E_x = -dV/dx$ ) and use this relation to solve problems   | Concept Check 3.7   | 77                               |
| 15  | Calculate the potential energy of a system of pair of charged particles  | FIGURE 3.30   | 79 & 80                          |
| <b>Important note: Please pay attention to specifying the units of measurement when solving problems, as grades will be calculated based on the units. Drawing relationships between variables, identifying and drawing the best fit line connecting the points, and finding values from the graph.</b> |  |   |                                  |
| Q1  | A<br>B<br>Develop a tool, equation or sketch, to obtain the resultant electric force exerted on a point charge by a nearby system of charges using the superposition principle   | EXERCISES 1.82 p: 25  | 10, 11,<br>& 12                  |
| Q2  | A<br>B<br>Develop a tool, equation or sketch, to obtain the resultant electric field strength at a point generated by a nearby system of point charges using the superposition principle<br>Solve problems related to the electric field due to several point charges  | As mentioned in the book  | 30 & 31                          |
| Q3  | A<br>B<br>Define the electric flux through a surface as the dot product between the electric field vector and the area vector at each point of that surface and expresses that in an equation (Solve problems on electric flux)<br>Prove that the electric flux through a closed surface is given by the net charge inside the surface divided by the permittivity of the medium, and write the Gauss's law in its integral form<br>(Apply Gauss' law to relate the net flux through a closed surface (real or imaginary) to the net charge enclosed by the surface) | EXAMPLE 2.5<br>As mentioned in the book   | 43<br>44, 45 & 46                |
| Q4  | A<br>B<br>Develop a mathematical equation to describe the electric potential of a point charge or many point charges or distributions of different charges   | As mentioned in the book FIGURE 3.21  | 70 & 71<br>79 & 80               |

**1** Apply Coulomb's law to relate the magnitude of the electrostatic force, the charge magnitudes of the pair if interacting particles, and the separation between them

**1. Electrostatic Force inside the Atom**

What is the magnitude of the electrostatic force that the two protons inside the nucleus of a helium atom exert on each other?

Where  $r = 2 \times 10^{-15} \text{ m}$  separates the two protons.

.....

.....

.....

.....

.....

What is the magnitude of the electrostatic force between a gold nucleus and an electron of the gold atom in an orbit with radius  $4.88 \times 10^{-12} \text{ m}$ ?

Where

the charge of the electron is  $qe = -e$

the charge of the gold nucleus is  $q_{\text{Nucleus}} = +79e$

.....

.....

.....

.....

**2. Equilibrium Position**

Two charged particles are placed as shown in Figure  $q_1 = 0.15 \mu\text{C}$  is located at the origin, and  $q_2 = 0.35 \mu\text{C}$  is located on the positive  $x$ -axis at  $x_2 = 0.40 \text{ m}$ .

Where should a third charged particle,  $q_3$ , be placed to be at an equilibrium point?

(such that the forces on it sum to zero)



.....

.....


.....

.....

3. Charged Balls


Two identical charged balls hang from the ceiling by insulated ropes of equal length,  $\ell = 1.50\text{m}$ . A charge  $q = 25.0\ \mu\text{C}$  is applied to each ball. Then the two balls hang at rest, and each supporting rope has an angle of  $25.0^\circ$  with respect to the vertical. What is the mass of each ball?

.....  
 .....  
 .....




Two balls have the same mass,  $0.9680\text{ kg}$ , and the same charge,  $29.59\ \mu\text{C}$ . They hang from the ceiling on strings of identical length,  $\ell$ , as shown in the figure. If the angle of the strings with respect to the vertical is  $29.79^\circ$ , What is the **length** of the strings?

.....  
 .....  
 .....



Two balls have the same mass and the same charge,  $15.71\ \mu\text{C}$ . They hang from the ceiling on strings of identical length,  $\ell = 1.223\text{ m}$ , As shown in the figure. The angle of the strings with respect to the vertical is  $21.07^\circ$ . What is the **mass** of each ball?

.....  
 .....  
 .....



Evaluate the **magnitude of the electrostatic** force exchanged between the two charges  $q_1 = +30\ \mu\text{C}$  and  $q_2 = -40\ \mu\text{C}$  separated by a distance of  $9.0\text{ cm}$

- (a)  $1.3 \times 10^3\text{N}$       (b)  $3 \times 10^3\text{ N}$       (c)  $3 \times 10^4\text{ N}$       (d)  $1.8 \times 10^3\text{N}$

The force between a charge of  $25\ \mu\text{C}$  and a charge of  $-10\ \mu\text{C}$  is  $8.0\text{ N}$ . What is the **separation between** the two charges?

- (a)  $0.28\text{ m}$       (b)  $0.53\text{ m}$       (c)  $0.45\text{ m}$       (d)  $0.15\text{ m}$

Two-point charges ( $q_1 = +q$ ) and ( $q_2 = -3q$ ) the distance between them is ( $25\text{ cm}$ ), if the electrostatic force between the two charges is ( $0.65\text{ N}$ ) What is the **value of second charge**?

- (a)  $1.2\text{ nC}$       (b)  $1.2\ \mu\text{C}$       (c)  $3.6\ \mu\text{C}$       (d)  $3.6\text{ nC}$


Two-point charges (+  $q$ ) and (-  $q$ ) have the same magnitude and the distance between them is (12 cm), if the electrostatic force between the two charges is (6.0 N) What is the value of each charge?

- (a) 8.9 nC      (b) 3.1  $\mu\text{C}$       (c) 8.9  $\mu\text{C}$       (d) 3.1 nC

Two-point charges (+  $q$ ) and (-  $q$ ) have the same magnitude and the distance between them is (9.0 cm), if the electrostatic force between the two charges is (5.0 N) What is the value of each charge?


- (a) 2.1 nC      (b) 2.1  $\mu\text{C}$       (c) 7.1 nC      (d) 7.1  $\mu\text{C}$

The figure shows two identical positively charged balls hanging from the ceiling by insulated massless ropes of equal length  $l$ . What is the charge on each ball?  
( $g=9.81 \text{ m/s}^2$ )




- (a) 9.1  $\mu\text{C}$       (c)  $9.3 \times 10^{-11} \mu\text{C}$   
(b) 9.6  $\mu\text{C}$       (d) 9.3  $\mu\text{C}$

The figure shows two identical positively charged balls hanging from the ceiling by insulated massless ropes of equal length  $l$ . What is the length  $l$ ?  
( $g=9.81 \text{ m/s}^2$ )



- (a) 8.28 cm      (c) 15.98 cm  
(b) 4.28 cm      (d) 0.68 m


Two identical positively charged balls hanging at rest from the ceiling by insulated massless ropes of equal length as shown in the figure. What is the distance between the two charges?



- (a) 3.0 m      (c) 0.5 m  
(b) 1.5 m      (d) 1.0 m

Two identical positively charged balls hanging at rest from the ceiling by insulated massless ropes of equal length as shown in the figure. What is the **mass** for each ball?


( $g=9.81 \text{ m/s}^2$ )



- (a) 0.2kg
- (b) 0.5kg
- (c) 0.6kg
- (d) 1.0kg


Two identical positively charged balls hanging at rest from the ceiling by insulated massless ropes of equal length as shown in the figure. What is the **mass** for each ball?

( $g=9.81 \text{ m/s}^2$ )



- (a) 0.2kg
- (b) 0.5kg
- (c) 0.6kg
- (d) 1.0kg



In the figure, the balls were given the same amount of charge, they repel and become to equilibrium as shown, the weight of each ball is  $1.20 \text{ N}$ , if  $r = 0.10 \text{ m}$ . What is the magnitude of the **charge** on each of them



- (a)  $1.0 \mu\text{C}$
- (b)  $1.2 \times 10^{-6} \text{ C}$
- (c)  $6.4 \times 10^{-7} \text{ C}$
- (d)  $0.72 \mu\text{C}$

Two identical charged balls hang from the ceiling by insulated ropes of equal length,  $\ell=0.8\text{m}$ .


A charge  $q=10.2\mu\text{C}$  is applied to each ball. Then the two balls hang at rest, and each supporting rope has an angle of  $24.0^\circ$  with respect to the vertical. What is the **mass** of each ball?



- (a) 0.51 kg
- (b) 0.57 kg
- (c) 0.67 kg
- (d) 0.77 kg




Two positive charges each of magnitude  $q$  are positioned on ( $y=+a$  and  $y=-a$ ) as shown where should we put a third charge  $Q$  on the  $y$ -axis such that the net force on the charge  $Q$  is zero.




- (a)  $y=0$       (b)  $y=-2a$       (c)  $y=+2a$       (d)  $y=\frac{a}{2}$

The figure shows two points charges, where can the electrostatic force be absent?



- (a) 1      (b) 2      (c) 3      (d) 4

The figure shows two points charges, where can the electrostatic force be absent?



- (a) 1      (b) 2      (c) 3      (d) 4

A charge  $Q_1$  is positioned on the  $x$ -axis at  $x = a$ . Where should a charge  $Q_2 = -4Q_1$  be placed to produce a net electrostatic force of zero on a third charge,  $Q_3 = Q_1$ , located at the origin?

- (a) at  $x = -a$       (b) at  $x = -2a$       (c) at  $x = 2a$       (d) at the origin

A charge  $Q_1 = Q$  is positioned on the  $x$  axis at  $x = a$ . Where should a charge  $Q_2 = 9Q$  be placed to produce a net electric force of zero on charge placed at the origin

- (a) at  $x = -3a$       (b) at  $x = -2a$       (c) at  $x = 2a$       (d) at  $x = 3a$

A point charge  $+3q$  is located at the origin, and a point charge  $-q$  is located on the  $x$ -axis at  $D = 0.500$  m. At what location on the  $x$ -axis will a third charge,  $q$ , experience no net force from the other two charges?

- (a) 0.68m      (b) 1.28m      (c) 1.18m      (d) 0.500m

Two-point charges are fixed on the  $x$ -axis:  $q_1 = 6.0 \mu\text{C}$  is located at the origin,  $O$ , with  $x_1 = 0.0$  cm, and  $q_2 = -3.0 \mu\text{C}$  is located at point  $A$ , with  $x_2 = 8.0$  cm. Where should a third charge,  $q_3$ , be placed on the  $x$ -axis so that the total electrostatic force acting on it is zero?




- (a) 19 cm      (b) 8.0 cm      (c) 0.0 cm      (d) 27 cm

2

Develop a tool, sketches, descriptive text or presentation to show the morphology of electric field lines of a single or multiple charge system with positive or negative charges


The spatial distribution of the electric field due to charges (1,2,3) is shown in the figure below Which of the parameters regarding the charges are correct?

|     | Positive Charge | Negative Charge | Magnitude of charges |
|-----|-----------------|-----------------|----------------------|
| (a) | A, C            | B               | $A > B > C$          |
| (b) | B               | A, C            | $B > A = C$          |
| (c) | B               | A, C            | $B > A > C$          |
| (d) | A, B, C         | None            | $B > A = C$          |




The spatial distribution of the electric field due to charges  $q_1$  and  $q_2$  is shown in the figure below.

|     | Charge $q_1$ | Charge $q_2$ | Magnitude   |
|-----|--------------|--------------|-------------|
| (a) | positive     | negative     | $q_1 > q_2$ |
| (b) | positive     | negative     | $q_2 > q_1$ |
| (c) | negative     | positive     | $q_1 > q_2$ |
| (d) | negative     | positive     | $q_2 > q_1$ |




The electric field lines for a system of two charges is shown below. Which of the following could be the correct charges 1 and 2?

|     | Charge $q_1$ | Charge $q_2$ |
|-----|--------------|--------------|
| (a) | $+32 \mu C$  | $-16 \mu C$  |
| (b) | $-32 \mu C$  | $+16 \mu C$  |
| (c) | $-16 \mu C$  | $+32 \mu C$  |
| (d) | $-32 \mu C$  | $-32 \mu C$  |




The figure shows three charges placed at three points A, B and C. Which of the following statements is correct about the charge (q) of A, B and C?

- (a)  $q_A = q_B$       (b)  $q_A = q_C$       (c)  $q_A = q_B = q_C$       (d)  $q_A = -q_C$



The figure shows three charges placed at three points A, B and C. Which of the following statements is incorrect about the charge?

- (a)  $q_A = q_C$       (b)  $q_B > -q_C$       (c)  $q_B > -q_A$       (d)  $q_A = -q_C$



3

Define the electric field vector at a point in space as the quotient of the electrostatic force vector acting on a positive test charge at this point divided by the test charge

Negative charge  $-2 \times 10^{-7} \text{ C}$  is located in an electric field of  $20 \text{ N/C}$  directed toward the East. What is the **electric force** acting on the charge?


- (a)  $4.0 \times 10^{-6} \text{ N East}$       (b)  $1.0 \times 10^{-8} \text{ N East}$       (c)  $4.0 \times 10^{-6} \text{ N West}$       (d)  $1.0 \times 10^{-8} \text{ N West}$

A uniform electric field of magnitude  $1000 \text{ N/C}$  is directed to the right from A to B. What would be the **magnitude and direction** of the electric force on a positive charge  $+20 \text{ nC}$  located close to point A?



- (a)  $2.0 \times 10^{-5} \text{ N A to B}$       (b)  $2.0 \times 10^{-10} \text{ N B to A}$       (c)  $2.0 \times 10^{-11} \text{ N A to B}$       (d)  $2.0 \times 10^{-11} \text{ N B to A}$

The figure shows two-point charges. if the electrostatic force attraction between them is  $6 \mu\text{N}$ . What is the **magnitude and direction of the electric field** acting on the positive charge?



- (a)  $2 \text{ N/C to the left}$       (b)  $2 \text{ N/C to the right}$       (c)  $6 \text{ N/C to the left}$       (d)  $6 \text{ N/C to the right}$

Which of the following represents force?

- (a)  $Eq$       (b)  $\frac{E}{q}$       (c)  $\frac{q}{E}$       (d)  $q^2E$

4

Solve problems involving general charge distribution and the electric field  
Find for a uniform distribution of charge, the linear charge density  $\lambda$  for charge along a line, the surface charge density  $\sigma$  for charge on a surface, and the volume

What is the **unit** of measuring the Linear charge density ( $\lambda$ ) on a thin metallic sheet?

- (a)  $\text{C/s}$       (b)  $\text{C/m}$       (c)  $\text{C/m}^2$       (d)  $\text{C/m}^3$

What is the **unit** of measuring the surface charge density ( $\sigma$ ) on a thin metallic sheet

- (a)  $\text{C/s}$       (b)  $\text{C/m}$       (c)  $\text{C/m}^2$       (d)  $\text{C/m}^3$

What is the **unit** of measuring the Volume charge density ( $\rho$ ) on a thin metallic sheet?

- (a)  $c/s$       (b)  $c/m$       (c)  $c/m^2$       (d)  $c/m^3$

What does  $x$  represent in the equation  $dq = X dV$  for a charge distribution over all the parts of an insulating sphere? And what **is the unit of  $x$** ?

- (a) *surface charge density  $C/m^2$*       (c) *linear charge density  $C/m$*   
 (b) *volume charge density  $C/m^3$*       (d) *infinity charge density  $C/m^4$*

If the charge is distributed over a **One- dimensional** object. What is the unit of charge density of this object?

- (a)  $c/s$       (b)  $c/m$       (c)  $c/m^2$       (d)  $c/m^3$

If the charge is distributed over a **two- dimensional** object. What is the unit of charge density of this object?

- (a)  $c/s$       (b)  $c/m$       (c)  $c/m^2$       (d)  $c/m^3$

If the charge is distributed over a **three- dimensional** object. What is the unit of charge density of this object?

- (a)  $c/s$       (b)  $c/m$       (c)  $c/m^2$       (d)  $c/m^3$

A long wire carries a charge  $12.0 \mu C/m$ , what is the **charge** of  $0.333 m$  of it?

- (a)  $1.2 \mu C$       (b)  $36 \mu C$       (c)  $1.8 \mu C$       (d)  $4.0 \mu C$

A conducting sphere has a charge  $23.5 nC$ , if its radius is  $25 cm$ , what its **charge surface density**?

Hint: Area of sphere ( $A = 4\pi r^2$ )

- (a)  $3 \times 10^{-8} C/m^2$       (b)  $6.0 \times 10^{-6} C/m^2$       (c)  $1.2 \times 10^{-6} C/m^2$       (d)  $4500 C/m^2$

A large, flat, horizontal sheet of charge has a charge per unit area of  $\sigma = 25.0 \mu C/m^2$ .

What is the **total charge** of the  $0.02 cm^2$  sheet?

- (a)  $0.5 \mu C$       (b)  $2.6 \mu C$       (c)  $25 pC$       (d)  $50 pC$

A 2D disk with a radius of  $12 cm$  has a uniform charge density of  $14 C/m^2$ .

What is the amount of **charge** distributed over the surface of this disk?

(Hint: Area of *disk*  $A = \pi r^2$ )

- (a)  $0.63 C$       (b)  $0.49 C$       (c)  $0.12 C$       (d)  $0.35 C$

A plastic disk with a diameter of 20 cm is charged uniformly with  $10^{14}$  electrons.  
What is its **surface charge density**

(Hint: Area of disk  $A = \pi r^2$ )

- (a)  $5.1 \times 10^{-6} \text{C/m}^2$       (b)  $1.3 \times 10^{-4} \text{C/m}^2$       (c)  $2.5 \times 10^{-4} \text{C/m}^2$       (d)  $5.1 \times 10^{-4} \text{C/m}^2$

Two long, uniformly charged rods are placed parallel to each other. Rod A has a total charge of  $10 \mu\text{C}$  and a length of 5.0 m. Rod B has a total charge of  $15 \mu\text{C}$  and a length of 3.0 m.

What is the **ratio** of the linear charge density  $\lambda_A$  of rod A to the linear charge density  $\lambda_B$  of rod B?

- (a)  $3/4$       (b)  $1/4$       (c)  $4/5$       (d)  $2/5$

A large plastic disc with a radius  $r$  is charged uniformly with  $4.5 \times 10^{18}$  electrons. If the surface charge density of this disc is  $0.11 \text{ C/m}^2$ , what is the **radius** of this disc?

- (a) 1.4 m      (b) 4.1 m      (c) 0.14 m      (d) 0.41 m

Given a linear charge density ( $\lambda = 5.0x^2$ ) from ( $x=0.0\text{m}$ ) to ( $x=4.0\text{m}$ ).  
Find **the electric charge** between ( $x=0.0\text{m}$ ,  $x=4.0\text{m}$ )

- (a) 106.6 C      (b) 40 C      (c) 0 C      (d) 853.3C

What does  $x$  represent in the formula  $E_y = \frac{2kx}{y}$  of an **infinitely long** wire and what is its unit?


- (a) *surface charge density C/m<sup>2</sup>*      (c) *linear charge density C/m*  
(b) *volume charge density C/m<sup>3</sup>*      (d) *infinity charge density C/m<sup>4</sup>*

**5** Apply the relationship between the electric field  $E$  and the electric force  $F$  and the charge  $q$

**Force due to an Electric Field**

An electron with a kinetic energy of 2.00 keV is fired horizontally across a horizontally oriented charged conducting plate with a surface charge density of  $+4.00 \times 10^{-6} \text{ C/m}^2$ . Taking the positive direction to be upward (away from the plate), What is the **vertical deflection** of the electron after it has traveled a horizontal distance of 4.00 cm? Where ( $1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$ )

$$y_f - y_0 = - \frac{e\sigma x_f^2}{2m\epsilon_0 v_0^2} = - \frac{e\sigma x_f^2}{4\epsilon_0 K}$$



A proton is placed in the uniform electric field of magnitude  $E = 0.25 \text{ N/C}$ . Find the **acceleration** of the proton (in  $\text{m/s}^2$ ). Hint: Proton mass is  $1.67 \times 10^{-27} \text{ kg}$  and proton charge is  $1.6 \times 10^{-19} \text{ C}$ .

- (a)  $2.5 \times 10^7$       (b)  $5.0 \times 10^8$       (c)  $6.0 \times 10^7$       (d)  $9.0 \times 10^{-7}$

An electron with velocity  $1.55 \times 10^3 \text{ m/s}$  is fired horizontally across a horizontally oriented charged conducting plate with a surface charge density of  $+3.0 \times 10^{-15} \text{ C/m}^2$ .

What is the **magnitude of vertical deflection** of the electron?

- (a)  $4.9 \times 10^{-3} \text{ m}$       (b)  $0.5 \text{ cm}$       (c)  $2.7 \times 10^{-6} \text{ m}$       (d)  $2.7 \text{ cm}$




As shown in the figure an electron is fired horizontally towards the positive x direction over a horizontally oriented charged conducting plate with a surface charge density of  $(+3.0 \times 10^{-15} \text{ C/m}^2)$ .

If the vertical deflection of the electron is  $(0.5 \text{ cm})$  after it has traveled a horizontal distance of  $(2.0 \text{ cm})$ .

What is the **kinetic energy** of the electron when is fired? (Neglect Earth gravity)


- (a)  $1.08 \times 10^{-24} \text{ J}$       (b)  $5.42 \times 10^{-24} \text{ J}$       (c)  $2.38 \times 10^6 \text{ J}$       (d)  $1.54 \times 10^3 \text{ J}$



As shown in the figure an electron is fired horizontally towards the positive x direction over a horizontally oriented charged conducting plate with a surface charge density of  $(+3.0 \times 10^{-15} \text{ C/m}^2)$ . If the vertical deflection of the electron is  $(0.5 \text{ cm})$  after it has traveled a horizontal distance of  $(2.0 \text{ cm})$ .

What is the **velocity** of the electron when is fired?

- (a)  $2.4 \times 10^6 \text{ m/s}$       (b)  $1.6 \times 10^3 \text{ m/s}$       (c)  $1.3 \times 10^5 \text{ m/s}$       (d)  $1.2 \times 10^3 \text{ m/s}$



According to the figure showing an electron fired with an initial velocity  $V_0$  from point A above a horizontally charged plate with a surface charge density of  $3.2 \mu\text{C/m}^2$ , the electron arrived at position (B).

What is the magnitude of  $V_0$  at point A

- (a)  $3.1 \times 10^{16} \text{ m/s}$       (b)  $3.5 \times 10^7 \text{ m/s}$       (c)  $3.0 \times 10^8 \text{ m/s}$       (d)  $1.8 \times 10^8 \text{ m/s}$




In the figure, a small negatively charged object is placed at rest in a uniform electric field. Which of the following statements **describes** the motion of the object when it is released? Neglect the mass



- (a) begin to move with a constant acceleration towards the right
- (b) begin to move with a constant speed towards the left
- (c) begin to move with an increasing acceleration towards the left.
- (d) begin to move with a constant acceleration towards the left

In the figure, a small positively charged object is placed at rest in a uniform electric field. Which of the following statements **describes** the motion of the object when it is released? Neglect the mass




- (a) begin to move with a constant acceleration towards the right
- (b) begin to move with a constant speed towards the left
- (c) begin to move with an increasing acceleration towards the left.
- (d) begin to move with a constant acceleration towards the left


6

Apply the relationship between the electric field  $E$  and the electric force  $F$  and the charge  $q$

The diagram shows a particle with positive charge  $Q$  and a particle with negative charge  $-Q$ . The electric field at point  $P$  on the perpendicular bisector of the line joining them is .....



The diagram shows two particles with positive charge  $+Q$ . The electric field at point  $P$  on the perpendicular bisector of the line joining them is .....




Consider two-point charges  $q_1 = +4.0\mu\text{C}$  and  $q_2 = -8.0\mu\text{C}$ , separated by a distance of 4.0 m. Find the magnitude of the electric field (in N/C) midway between the two-point charges

- (a)  $2.7 \times 10^4$       (b)  $72 \times 10^4$       (c)  $9.0 \times 10^3$       (d)  $1.8 \times 10^4$


According to the figure, two charges ( $Q = -3.0\mu\text{C}$ ) and ( $q = +2.0\mu\text{C}$ ), the electric force between them equals to  $7.3 \times 10^{-4}$  N and the distance between them is (8.6m), if q is placed at (5m,0m).

What is the electric field at the point (0,0)?




- (a) 1271.0N/C      (b) 905.0N/C      (c) -551.0N/C      (d) 720.02N/C

According to the figure, two charges ( $Q = +8.0\mu\text{C}$ ) and ( $q = -5.0\mu\text{C}$ ), the electric force between them equals to  $(3.6 \times 10^{-1}$  N), if Q is placed at (0 m, 0.8m). What is the electric field at the point (0,0)?



- (a)  $1.68 \times 10^5$  N/C      (b)  $1.12 \times 10^5$  N/C      (c)  $1.25 \times 10^5$  N/C      (d)  $2.82 \times 10^{10}$  N/C


1. According to the figure, if ( $Q = -2.0\mu\text{C}$ ) and ( $q = +2.0\mu\text{C}$ ), what is the electric field at the point P(0,0)



- (a)  (b) 
- (c)  (d) 



The figure shows the electric field pattern around two charges of equal magnitudes and opposite signs. Which of the labeled arrows correctly represents the direction of the electric field vector at point P?




- (a) A                      (b) B                      (c) C                      (d) D

9

Solve problems on electric flux Define the electric flux through a surface as the dot product between the electric field vector and the area vector at each point of that surface and expresses that in an equation

A cubical Gaussian surface is placed in a uniform electric field as shown in the figure. The length of each edge of the cube is 1.0 m. The uniform electric field has a magnitude of  $5.0 \times 10^8 \text{ N/C}$  and passes through the left and right sides of the cube perpendicular to the surface. What is the total electric flux that passes through the cubical Gaussian surface?



- (a) zero                      (b)  $3.0 \times 10^9 \text{ Nm}^2/\text{C}$                       (c)  $1.5 \times 10^7 \text{ Nm}^2/\text{C}$                       (d)  $2.5 \times 10^6 \text{ Nm}^2/\text{C}$

A flat surface of area  $3.20 \text{ m}^2$  is rotated in a uniform electric field of magnitude  $E = 6.20 \times 10^2 \text{ N/C}$ . Determine the electric flux through this area when the electric field is perpendicular to the surface


- (a)  $0 \text{ Nm}^2/\text{C}$                       (b)  $1.98 \times 10^6 \text{ Nm}^2/\text{C}$                       (c)  $1.40 \times 10^6 \text{ Nm}^2/\text{C}$                       (d)  $6.19 \times 10^5 \text{ Nm}^2/\text{C}$

A flat surface of area  $3.20 \text{ m}^2$  is rotated in a uniform electric field of magnitude  $E = 6.20 \times 10^2 \text{ N/C}$ . Determine the electric flux through this area when the electric field is parallel to the surface

- (a)  $0 \text{ Nm}^2/\text{C}$                       (b)  $1.98 \times 10^6 \text{ Nm}^2/\text{C}$                       (c)  $1.40 \times 10^6 \text{ Nm}^2/\text{C}$                       (d)  $6.19 \times 10^5 \text{ Nm}^2/\text{C}$


According to the figure, a cube that has (5.0cm) side length in a uniform electric field ( $E = 200 \text{ N/C}$ ), that is perpendicular to the plane of one face of the cube.

What Is the magnitude of electric flux passing through the black face?




- (a)  $0 \text{ Nm}^2/\text{C}$                       (b)  $1.0 \text{ Nm}^2/\text{C}$                       (c)  $1.5 \text{ Nm}^2/\text{C}$                       (d)  $0.5 \text{ Nm}^2/\text{C}$

According to the figure, a uniform electric field ( $E = 360\text{N/C}$ ), that is perpendicular to the plane of one face of the cube. if the electric flux passing through the left shaded face is equal to  $(-1.2\text{Nm}^2/\text{C})$ . What is the cube side length?




- (a) 0.058m      (b)  $3.3 \times 10^{-3}\text{m}$       (c) 17.3m      (d) 300m

According to the figure, a uniform electric field ( $E = 28\text{N/C}$ ), that is perpendicular to the plane of one face of the cube. If the electric flux passing through the left shaded face is equal to  $(-7.0\text{Nm}^2/\text{C})$ , what is the volume the cube?




- (a)  $0.125\text{m}^3$       (b)  $8.000\text{m}^3$       (c)  $0.250\text{m}^3$       (d)  $0.500\text{m}^3$

What is the flux through a circular area with radius of  $0.30\text{m}$  placed in an external electric field has a strength of  $1200\text{N/C}$  and makes an angle of  $60^\circ$  with the plane of the circle?




- (a)  $293\text{Nm}^2/\text{C}$       (b)  $169\text{Nm}^2/\text{C}$       (c)  $565\text{Nm}^2/\text{C}$       (d)  $979\text{Nm}^2/\text{C}$

In the figure a cube with sides of length  $5.0\text{cm}$ , an electric field is passes through the cube if the electric field is given by this equation:  $E = 2.0x\hat{i} + 4.0y\hat{j} + 6.0z\hat{k}$ . What is the electric flux through the shaded faces?




- (a)  $0.015\text{Nm}^2/\text{C}$       (b)  $0.020\text{Nm}^2/\text{C}$       (c)  $0.030\text{Nm}^2/\text{C}$       (d)  $0.0050\text{Nm}^2/\text{C}$

According to the figure, at which ( $\theta$ ) the magnetic flux equal approximately to  $(0.5EA)$ ?




- (a) 60      (b) 30      (c) 90      (d) 0

According to the figure, which of the following is **corresponding** about the electric flux from surface A?



- (a)  $\frac{\phi_E}{\phi_{max}} = \frac{3}{5}$       (b)  $\frac{\phi_E}{\phi_{max}} = \frac{4}{5}$       (c)  $\frac{\phi_E}{\phi_{max}} = \frac{5}{3}$       (d)  $\frac{\phi_E}{\phi_{max}} = \frac{5}{4}$

The diagram shows a ring in a uniform electric field passing through its surface. The ring rotates so that the angle it makes with the field changes as shown in the diagram, the angle made by the ring from position (a) to position (b) is 17, what is **the electric flux  $\phi_b$** ?




- (a)  $9.60 Nm^2/C$       (b)  $16.8 Nm^2/C$       (c)  $15.7 Nm^2/C$       (d)  $12.2 Nm^2/C$

10

Prove that the electric flux through a closed surface is given by the net charge inside the surface divided by the permittivity of the medium, and write the Gauss's law in its integral form


$$\oiint \vec{E} \cdot d\vec{A} = \frac{q}{\epsilon_0}$$

Three isolated charges of +2q, -2q, and +3q are placed in a 3D vacuum space, where they are surrounded by a Gaussian surface, as shown in the figure. What is **the total electrical flux** through that surface?



- (a)  $\phi = \frac{+7q}{\epsilon_0}$       (b)  $\phi = \frac{+3q}{\epsilon_0}$       (c)  $\phi = +3q$       (d)  $\phi = \frac{+5q}{4\pi\epsilon_0}$

Assume that the cube shown in the figure contains seven electrons, eight neutrons, and a number of protons, if the electric flux through the cube is  $(3.62 \times 10^{-8} \text{ Nm}^2/\text{C})$ . How many protons in the cube?




- (a) 7                      (b) 8                      (c) 9                      (d) 2

what is the magnitude of the electric flux through the sphere shown in the figure that contains an electron and two protons?



- (a)  $1.6 \times 10^{-8} \text{ Nm}^2/\text{C}$       (b)  $1.8 \times 10^{-8} \text{ Nm}^2/\text{C}$       (c)  $3.6 \times 10^{-8} \text{ Nm}^2/\text{C}$       (d)  $5.4 \times 10^{-8} \text{ Nm}^2/\text{C}$

A neutral sphere, made of insulating material, is placed in an external electric field as shown in the figure. The net electrical flux passing through the surface of the sphere is:




- (a) Zero                      (b) Negative                      (c) positive                      (d) cannot be determined

A small sphere has a charge  $+q$ . Spherical Gaussian Surfaces A and B are concentric with the sphere, as shown in the figure below. The radii of surfaces A and B are  $r$  and  $2r$ , respectively.

The magnitude of the electric flux through A is  $\phi_A$ . The magnitude of the electric flux through surface B is  $\phi_B$ .

The ratio  $\frac{\phi_A}{\phi_B}$  is



Hint: Area of sphere ( $A = 4\pi r^2$ )

- (a)  $\frac{4}{1}$                       (b)  $\frac{2}{1}$                       (c)  $\frac{1}{1}$                       (d)  $\frac{1}{2}$

A point charge sits in the center of a 1 m cube. The electric flux through one side of the cube is  $7.2 \times 10^{10} \text{ Nm}^2/\text{C}$ . What is the charge at the center of the cube?




- (a) 3.8 C                      (b) 4.6 C                      (c) 6.0 C                      (d) 8.9 C

A hollow spherical conductor with an inner radius 5.0 cm and an outer radius 6.5 cm and its outer surface carries an electric charge, an electric charge  $q$  is placed at its center, resulting in an electric field at the inner surface of the conductor  $50 \text{ N/C}$  towards the center of the conductor and another electric field at the outer surface of the conductor  $50 \text{ N/C}$  away from the center of the conductor, what is the **magnitude and type of charge  $q$** ?


- (a)  $+2.5 \times 10^{-12} \text{ C}$       (b)  $-2.5 \times 10^{-12} \text{ C}$       (c)  $-1.4 \times 10^{-11} \text{ C}$       (d)  $+1.4 \times 10^{-11} \text{ C}$

The figure below shows five Gaussian surfaces (a to e) surrounding a distribution of charges. Which of the Gaussian surfaces have **the largest electric flux**



- (a) a      (b) b      (c) c      (d) d

The figure below shows a distribution of charges. The **flux of the electric** field due to these charges through the surface S is



- (a)  $\phi = \frac{+3q}{\epsilon_0}$       (b)  $\phi = \frac{+2q}{\epsilon_0}$       (c)  $\phi = +3q$       (d)  $\phi = \frac{+5q}{4\pi\epsilon_0}$

Inside a spherical surface is  $5.3 \times 10^{-6} \text{ C}$  and a  $-2.2 \times 10^{-6} \text{ C}$  charge. What is **the total electric flux** through the surface of the sphere in units of  $\text{Nm}^2/\text{C}$ ?

- (a)  $3.4 \times 10^{-16}$       (b)  $3.1 \times 10^6$       (c)  $3.5 \times 10^5$       (d)  $2.8 \times 10^4$

A  $3.5 \text{ C}$  point charge sits in the center of a  $1 \text{ m}$  cube. What is the **electric flux** through one side of the cube?

- (a)  $6.6 \times 10^{10} \text{ Nm}^2/\text{C}$       (b)  $4.5 \times 10^{10} \text{ Nm}^2/\text{C}$       (c)  $3.3 \times 10^{10} \text{ Nm}^2/\text{C}$       (d)  $5.0 \times 10^{10} \text{ Nm}^2/\text{C}$

11

Prove that the electric flux through a closed surface is given by the net charge inside the surface divided by the permittivity of the medium, and write the Gauss's law in its integral form

The magnitude of the electric field due to an infinite, flat, thin and nonconducting plane of charge is  $(1.55 \times 10^4 \text{ N/C})$ . Assuming that the charge is uniformly distributed, What is **the charge density** on the surface of the plane?


- (a)  $3.10 \times 10^{-8} \text{ C/m}^2$     (b)  $1.55 \times 10^{-8} \text{ C/m}^2$     (c)  $1.37 \times 10^{-7} \text{ C/m}^2$     (d)  $2.74 \times 10^{-7} \text{ C/m}^2$

The diagram below shows two charged plates placed close to each other. Rank points 1, 2 and 3 from greatest to the least electric field.




- (a)  $1 > 2 > 3$     (b)  $1 = 2 = 3$     (c)  $3 = 2 > 1$     (d)  $3 > 1 > 2$

In the diagram below, two infinitely thin, parallel, nonconducting plates are placed in the air, separated by a distance of  $d$ . They each carry a uniform positive charge with density distribution of  $\sigma$ . What is **the magnitude of the electric field** at point aa



- (a)  $\frac{\sigma}{2\epsilon}$     (b)  $\frac{\sigma}{\epsilon}$     (c)  $\frac{2\sigma}{\epsilon}$     (d) 0

In the diagram below, two infinitely thin, parallel, nonconducting plates are placed in the air, separated by a distance of  $d$ . They each carry a uniform positive and negative charge with density distribution of  $\sigma$ . What is **the magnitude of the electric field** at point aa



- (a)  $\frac{\sigma}{2\epsilon}$     (b)  $\frac{\sigma}{\epsilon}$     (c)  $\frac{2\sigma}{\epsilon}$     (d) 0

The magnitude of the electric field due to an infinite, flat, thin and conducting plane of charge is  $(1.55 \times 10^4 \text{ N/C})$ . Assuming that the charge is uniformly distributed, What is **the charge density** on the surface of the plane?

مقدار المجال الكهربائي الناتج عن لوح مسطح رقيق لانتهائي و موصل، مشحون بشحنة كهربائية هو  $(1.55 \times 10^4 \text{ N/C})$ . بافتراض أن الشحنة موزعة بانتظام، ما كثافة الشحنة على السطح؟

- (a)  $3.10 \times 10^{-8} \text{ C/m}^2$     (b)  $1.55 \times 10^{-8} \text{ C/m}^2$     (c)  $1.37 \times 10^{-7} \text{ C/m}^2$     (d)  $2.74 \times 10^{-7} \text{ C/m}^2$

As shown in the figure. **nonconducting surface**

Find the total electric field (**magnitude and direction**) at each ( $A - B - C$ )

كما هو موضح في الشكل. سطح غير موصل. أوجد المجال الكهربائي الكلي (المقدار والاتجاه) عند كل من ( $A - B - C$ )



## 12

## Solve problems involving electric potential energy

Which of the following statements is **correct**?

- (a) The change in electric potential energy due to some spatial rearrangement of a system is equal to the negative of the work done by the conservative force during this spatial rearrangement
- (b) The change in electric potential energy due to some spatial rearrangement of a system is equal to the positive of the work done by the conservative force during this spatial rearrangement
- (c) The change in electric potential energy due to some spatial rearrangement of a system is equal to the positive of the work done by the unconservative force during this spatial rearrangement
- (d) The change in electric potential energy due to some spatial rearrangement of a system is equal to the negative of the work done by the unconservative force during this spatial rearrangement.


For a **proton** moving in the direction of the electric field

- (a) its potential energy increases and its electric potential decreases.
- (b) its potential energy decreases and its electric potential decreases.
- (c) its potential energy increases and its electric potential increases.
- (d) its potential energy decreases and its electric potential increases.

For an **electron** moving in a direction opposite to the electric field


- (a) its potential energy increases and its electric potential decreases.      (c) its potential energy increases and its electric potential increases.
- (b) its potential energy decreases and its electric potential decreases.      (d) its potential energy decreases and its electric potential increases.

A positive charge of  $3.0 \times 10^{-8} \text{ C}$  is placed in an upward directed uniform electric field of  $4.0 \times 10^4 \text{ N/C}$ . When the charge is moved 0.5 m upward, the **work done by** the electric force on the charge is:



- (a)  $6 \times 10^{-4} \text{ J}$       (c)  $8 \times 10^4 \text{ J}$
- (b)  $12 \times 10^{-4} \text{ J}$       (d)  $2 \times 10^4 \text{ J}$

A proton is released from rest in  $300.0 \text{ N/C}$  electric field pointing to positive x-direction. Calculate the **change in electric potential energy** if it moved 10.0 cm making an angle  $60.0^\circ$  with the electric field.



- (a)  $-2.40 \times 10^{-18} \text{ J}$       (c)  $2.40 \times 10^{-16} \text{ J}$
- (b)  $2.40 \times 10^{-18} \text{ J}$       (d)  $2.40 \times 10^{-16} \text{ J}$

A proton is accelerated from rest close to the positive plate to deliver to the negative plate with maximum kinetic energy  $4.8 \times 10^{-17} \text{ J}$

What is the **absolute value of electric potential** difference between these two parallel plates?

- (a) 0V      (c) 3V
- (b) 30V      (d) 300V

A proton, initially at rest, is accelerated through an electric potential difference of 500 V. What is the **kinetic energy** of the proton?

- (a) 500 J      (b)  $+1.6 \times 10^{-19} \text{ J}$       (c)  $+8.0 \times 10^{-17} \text{ J}$       (d) zero

An electron is accelerated from rest close to the negative plate of a capacitor. It reaches the positive plate in (0.02s). If the electric potential difference between these two plates is (100V)?

What is the **acceleration** of the electron?

- (a)  $2.96 \times 10^8 \text{ m/s}^2$       (b)  $6.92 \times 10^6 \text{ m/s}^2$       (c)  $1.92 \times 10^{10} \text{ m/s}^2$       (d)  $3.80 \times 10^{10} \text{ m/s}^2$




An electron is accelerated from rest close to the negative plate to deliver to the positive plate, if the electric potential difference between the two parallel plates is (120V).  
What is the maximum kinetic energy of the electron?

- (a)  $1.9 \times 10^{-17} \text{ J}$       (b)  $1.6 \times 10^{-19} \text{ J}$       (c)  $9.6 \times 10^{-18} \text{ J}$       (d)  $3.8 \times 10^{-18} \text{ J}$

**13** Develop a method such as schematic representations to compare the equipotential surfaces due to a point charge, two identical charges, and two different charges

The figure shows the electric field lines and equipotential surfaces of two-point charges. Which of the following is **correct**?






- | point charges                        | J lines                |
|--------------------------------------|------------------------|
| (a) same magnitude and opposite sign | electric field lines   |
| (b) same magnitude and opposite sign | Equipotential surfaces |
| (c) identical and positive           | electric field lines   |
| (d) Equipotential surfaces           | identical and negative |

Which one of the following statements is **not** true?

- (a) Equipotential lines are parallel to the electric field lines.      (c) Equipotential surfaces exist for any charge distribution.  
(b) When a charge moves on an equipotential surface, the work done on the charge is zero.      (d) Equipotential lines for a point charge are circular.

The figure shows the electric field lines and equipotential surfaces of two-point charges. Which of the following is **correct**?

- | point charges              | red lines              |
|----------------------------|------------------------|
| (a) identical and positive | electric field lines   |
| (b) identical and positive | equipotential surfaces |
| (c) identical and negative | electric field lines   |
| (d) identical and negative | equipotential surfaces |



Which of the following **is/are equipotential surface/s**?

- (a) A      (b) B      (c) C      (d) A&B

Which of the following is **not equipotential surface/s**?

- (a) A      (b) B      (c) C      (d) A&B

Determine the **wrong** statement in the following.

- (a) we need to do work on a charge to move it on an equipotential surface, (c) The surface of any conductor is an equipotential surface.  
 (b) Electric field lines are perpendicular to equipotential surfaces at any point. (d) In a uniform electric field, the electric field lines are always parallel.

## 14

Relate the component of the electric field along a certain direction  $E_s$  to the change in the electric potential along that direction ( $E_s = -\frac{dV}{ds}$ ) and use this relation to solve problems

Suppose that the voltage at a point is given by the equation ( $V_{(x,y,z)} = 8x^2 - 9y^2 + 5z^2$ ) in volts. Which of the **dimensions** (x, y, z) determines the magnitude of the electric field at this point?

- (a) x (b) y (c) z (d) x&y&z

Suppose that the voltage at a point is given by the equation ( $V_{(x,y,z)} = 8x - 9y + 5z^2$ ) in volts. Which of the **dimensions** (x, y, z) determines the magnitude of the electric field at this point?

- (a) x (b) y (c) z (d) x&y&z

Suppose an electric potential has the equation  $V_{(x,y,z)} = 3x - 6y + 2z$  in volts. What is the **magnitude** of associated electric field, in units of volts per meter at P (0,0,0)?

- (a) 0 (b) 7 (c) 1 (d) 6

The electric potential in some region is given by  $V_{(x,y)} = 2x^2 - 3y$ . Find the **x component of the electric field** associated with this potential at point (1,2)

- (a) 5 V/m (b) -6 V/m (c) -4 V/m (d) 8 V/m

The electric potential in some region is given by  $V_{(x,y)} = 3x - 2y^2$ . Find the **Y component of the electric field** associated with this potential at point (1,2)

- (a) 5 V/m (b) -6 V/m (c) -4 V/m (d) 8 V/m

An electric potential is described in volts by  $V(x, y, z) = 3x^2 + 8y - 6z$ . What is the **magnitude of an electric field** at the point (+2.0 m, -2.0 m, -1.0 m)

- (a) 27.6 V/m (b) 31.0 V/m (c) 15.6 V/m (d) 14.0 V/m


Suppose the electric potential is given by the equation  $V(x, y, z) = 5x^2 - 8y^2$   
What is the **magnitude of electric field** at the point (3,5,2)?

- (a) 7300 V/m (b) 10 V/m (c) 50 V/m (d) 85 V/m

15


Calculate the potential energy of a system of pair of charged particles

Consider two identical charges of  $q = 50\mu\text{C}$  each, placed 5.0 m apart.  
Find the **electrostatic potential energy stored** in the configuration.




- (a) 4.5J (b) 0.9J (c) 0.2J (d) 2.5J

The potential energy of a system of three equal charges arranged in an equilateral triangle is 0.54 J  
If the length of one side of this triangle is 33 cm,  
What is the **charge** of one of the three charges?



- (a) 1.7  $\mu\text{C}$  (b) 4.3  $\mu\text{C}$  (c) 2.6  $\mu\text{C}$  (d) 2.0  $\mu\text{C}$

Two ( $+ 4.0\mu\text{C}$ ) point charges arranged at the corners of a square with side length (12 cm)  
What is the potential energy of a system of two charges?



- (a) 1.2 J (b) 0.85 J (c) 10 J (d) 5.0 J

What happens to the **magnitude of electric potential energy** of a system of two-point charges when the distance between them decreases?

- (a) increases (b) decreases (c) remains the same (d) increases rapidly then decreases.

When two charges are separated by a distance  $d$ , their electric potential energy is equal to  $U$ .  
What would be their electric potential energy if the separation distance was  $\frac{d}{2}$ ?

- (a)  $U/4$  (b)  $U/2$  (c)  $2U$  (d)  $4U$

The electric potential energy of a system of two negative point charges is equal to  $1.2 \times 10^{-2}J$ , if the distance between them becomes four times what it was, what is **the potential energy** of the system?

- (a)  $1.5 \times 10^{-2}J$       (b)  $1.2 \times 10^{-2}J$       (c)  $4.8 \times 10^{-2}J$       (d)  $3.0 \times 10^{-3}J$

If the electric potential energy of a system of two negative point charges is (8.0J). What is the **electric potential energy** of the system when the distance between the two charges is halved?

- (a) 0.0J      (b) 4.0J      (c) 8.0J      (d) 16.0J

If the electric potential energy of a system of two negative point charges is (24.0J). What is the **electric potential energy** of the system when the distance between the two charges is tripled?

- (a) 0.0J      (b) 4.0J      (c) 8.0J      (d) 16.0J

الحمد لله رب العالمين تم الانتهاء من الجزء الإلكتروني

[https://t.me/EINSTEIN\\_AE](https://t.me/EINSTEIN_AE)

## PAPAR PART

Three-point charges are placed at the vertices of a right triangle as in the adjacent figure, If

The air surrounded the charges,  $q_1 = 6.0 \mu\text{C}$  and  $q_2 = 8.0 \mu\text{C}$   $q_3 = -q_2$

توضع الشحنات ثلاثية النقاط عند رؤوس المثلث القائم الزاوية كما في الشكل المجاور، أطاق الهواء بالشحنات،

Calculate the magnitude and direction of the electric force for charge  $q_3$

احسب مقدار القوة الكهربائية واتجاهها للشحنة  $q_3$

.....

.....

.....


.....

.....

.....

.....

.....



A charge of  $+10 \mu\text{C}$  is placed at the origin. Two other charges of  $-20 \mu\text{C}$  each are placed at equal distances from the origin as shown below.

شحنة  $+10 \mu\text{C}$  عند نقطة الأصل. وُضعت شحنتان أخريتن كل منهما  $-20 \mu\text{C}$  كل منهما عند مسافة متساوية من نقطة الأصل، كما هو موضح أدناه.

Calculate the magnitude and the direction of the net electric force on the  $+10 \mu\text{C}$  charge

احسب مقدار واتجاه القوة الكهربائية المحصلة المؤثرة على الشحنة  $+10 \mu\text{C}$

.....

.....

.....


.....

.....

.....

.....

.....



Three-point charges are placed as shown in the figure:

توضع الشحنات ثلاثية النقاط كما هو موضح في الشكل:


A- Calculate the magnitude of the resultant electrostatic force acting on the charge  $-7.0 \mu\text{C}$ .

احسب مقدار القوة الكهروستاتيكية المحصلة المؤثرة على الشحنة  $-7.0 \mu\text{C}$ .

B- Find the direction of the resultant electrostatic force acting on the charge  $-7.0 \mu\text{C}$ .


(By finding the angle)

أوجد اتجاه القوة الكهروستاتيكية المحصلة المؤثرة على الشحنة  $-7.0 \mu\text{C}$  (الزاوية بإيجاد)



The charges  $q_1 = 2.00 \times 10^{-7}\text{C}$ ,  $q_2 = -4.00 \times 10^{-7}\text{C}$ , and  $q_3 = -1.00 \times 10^{-7}\text{C}$ , are placed at the corners of the triangle shown below. What is the net force on  $q_1$ ?

الشحنات  $q_1 = 2.00 \times 10^{-7}\text{C}$ ,  $q_2 = -4.00 \times 10^{-7}\text{C}$ , and  $q_3 = -1.00 \times 10^{-7}\text{C}$ ، موضوعة عند زوايا المثلث الموضح بالأسفل. ما القوة المحصلة المؤثرة على  $q_1$ ؟



Two ( $-5.0\mu\text{C}$ ) fixed point charges are located in  $x y$  plane at points

$$(x = -1.0 \text{ cm}, y = +0.0 \text{ cm}), (x = +8.0 \text{ cm}, y = 0.0 \text{ cm})$$

Calculate the magnitude of electrostatic force acting on a charge of ( $+5.0 \mu\text{C}$ ) placed at ( $x = +8.0 \text{ cm}, y = +5.0 \text{ cm}$ ). And calculate the angle that the electrostatic force vector makes with the  $x$ -axis.

شحنتان نقطيتان متماثلتان كل منها ( $-5.0\mu\text{C}$ ) في المستوى ( $x y$ ) عند المواقع الآتية:

$$(x = -1.0 \text{ cm}, y = +0.0 \text{ cm}), (x = +8.0 \text{ cm}, y = 0.0 \text{ cm})$$

وضعت شحنة ثالثة ( $+5.0 \mu\text{C}$ ) عند نقطة موقعها ( $x = +8.0 \text{ cm}, y = +5.0 \text{ cm}$ ).

احسب مقدار القوة الكهروستاتيكية المؤثرة في الشحنة ( $+5.0 \mu\text{C}$ ) والزاوية التي يصنعها متجه القوة الكهروستاتيكية مع المحور  $x$  الموجب.


Find the net force on a  $+2.00\text{-C}$  charge at the origin of an  $xy$ -coordinate system if there is a  $+5.00 \text{ C}$  charge at  $(3.00 \text{ m}, 0.00)$  and a  $-3.00 \text{ C}$  charge at  $(0.00, 4.00 \text{ m})$ .

أوجد محصلة القوى المؤثرة في شحنة  $+2.00\text{-C}$  عند نقطة الأصل في نظام إحداثي  $xy$  إذا كانت هناك شحنة  $+5.00\text{-C}$  عند النقطة  $(3.00 \text{ m}, 0.00)$  وشحنة  $-3.00\text{-C}$  عند النقطة  $(0.00, 4.00 \text{ m})$ .

Three charges ( $q_3$ ,  $q_2$ , and  $q_1$ ) are placed in a space as shown in the figure below If ( $q_1 = -4\mu\text{C}$ ), ( $q_2 = 8\mu\text{C}$ ), and ( $q_3 = 6\mu\text{C}$ ) find the magnitude of the electric force acting on the charge  $q_2$


If charge ( $q_3$ ) is removed from charge ( $q_2$ ) and charge ( $q_1$ ) remains in place, does the magnitude of the electric force on ( $q_2$ ) increase, decrease, or remain constant and why?

وضعت ثلاث الشحنات ( $q_3$ ,  $q_2$ ,  $q_1$ ) في الفراغ كما هو مبين في الشكل المجاور  
إذا كانت ( $q_1 = -4\mu\text{C}$ ), ( $q_2 = 8\mu\text{C}$ ), ( $q_3 = 6\mu\text{C}$ ) اوجد مقدار القوة الكهربائية المؤثرة في الشحنة  $q_2$   
إذا أبعدت الشحنة ( $q_3$ ) نهائيا عن الشحنة ( $q_2$ ) مع بقاء ( $q_1$ ) في مكانها  
فهل يزداد مقدار القوة الكهربائية المؤثرة في ( $q_2$ ) أم يقل أم يبقى ثابتا ولماذا ؟



In the figures below, three-point charges are placed in a space If the net electric force on charge ( $q_1$ ) is zero where ( $q_2 = +4\mu\text{C}$ ) and ( $q_1 = -2\mu\text{C}$ ), find the magnitude of charge  $q_3$ .

في الشكل المجاور ثلاث الشحنات النقطية موضوعة في الفراغ إذا كانت محصلة القوى الكهربائية في الشحنة ( $q_1$ ) تساوي صفرا علما بان ( $q_1 = -2\mu\text{C}$ ), ( $q_2 = +4\mu\text{C}$ ) اوجد مقدار الشحنة  $q_3$






Four identical charged particles ( $q = +10.0 \mu\text{C}$ ) are located on the corners of a rectangle as shown in Figure. The dimensions of the rectangle are  $L=60.0 \text{ cm}$  and  $W= 15.0 \text{ cm}$ .


Calculate the **magnitude and the direction** of the total electric force exerted on the charge at **the lower left corner** by the other three charges.

أربعة جسيمات مشحونة متطابقة ( $q = +10.0 \mu\text{C}$ ) تقع على زوايا مستطيل كما هو موضح في الشكل. أبعاد المستطيل  $L=60.0 \text{ cm}$  و  $W= 15.0 \text{ cm}$ . احسب مقدار واتجاه القوة الكهربائية الكلية المبذولة على الشحنة في الزاوية اليسرى السفلية بواسطة الشحنات الثلاث الأخرى.



In the figure, the net electrostatic force on charge  $Q_A$  is zero. If  $Q_A = +1.00 \text{ nC}$ , determine the **magnitude** of  $Q_0$ .

في الشكل، القوة الكهروستاتيكية المحصلة المؤثرة على الشحنة  $Q_A$  تساوي صفراً. إذا كانت  $Q_A = +1.00 \text{ nC}$ ، فأوجد مقدار  $Q_0$ .




The diagram below is the electric field vector resulting at point p, which is in the path of two-point charges. If air is around the two charges and point p, what is the type of charge  $q_2$ ,  $q_1$ ?

Find the magnitude of the electric field acting on charge  $q_2$  if  $|q_1| = 3nC$

يبين الشكل المجاور متجه شدة المجال الكهربائي المحصلة عند النقطة (p) والواقعة في مجال شحنتين نقطيتين إذا كان الهواء يحيط بالشحنتين والنقطة (p)

ما نوع كل من الشحنتين  $q_2$ ,  $q_1$ ؟

أوجد مقدار شدة المجال الكهربائي المؤثر في الشحنة ( $q_2$ ) إذا كانت  $|q_1| = 3nC$



يوضح الشكل شحنتان نقطيتان ( $q_1 = -4\mu C$ ), ( $q_2 = +16\mu C$ )

أصعب مقدار شدة المجال الكهربائي عند نقطة أصل؟

أصعب مقدار القوة الكهربائية المؤثرة في الشحنة ( $q_1$ ) وحدد اتجاهها على الرسم.


إذا أزيلت الشحنة ( $q_2$ ) فهل يزداد مقدار المجال الكهربائي عند نقطة الأصل أم يقل أم لا يتغير؟ فسر اجابتك

The diagram below two-point charges ( $q_1 = -4\mu C$ ), ( $q_2 = +16\mu C$ )

Calculate the magnitude of the electric field at the origin.


Calculate the magnitude of the electric force acting on charge ( $q_1$ ) and draw direction

If charge ( $q_2$ ) is removed, does the magnitude of the electric field at the origin increase, decrease, or remain unchanged? Explain your answer



The figure shows the magnitude and the direction of electric field acting on a positive test charge You place at point O near two charges points  
Calculate the **magnitude of the charge and determine its type?**

يظهر الشكل المجاور مقدار واتجاه شدة المجال الكهربائي المؤثر في شحنة اختبار موجبة وضعت عند النقطة O بالقرب من شحنتين نقطيتين ، احسب مقدار الشحنة و حدد نوعها ؟



Two identical point charges each ( $-6 \text{ nC}$ ) are placed in the  $x, y$  plane at the following locations:

$(0.0 \text{ cm}, 0.0 \text{ cm})$  and  $(+3.0 \text{ cm}, +3.0 \text{ cm})$ .

Calculate the magnitude and direction of the resultant electric field at  $(0.0 \text{ cm}, +3.0 \text{ cm})$ .

شحنتان نقطيتان متماثلتان كل منهما ( $-6 \text{ nC}$ ) وضعت في المستوى  $x, y$  عند المواقع الآتية:


$(0.0 \text{ cm}, 0.0 \text{ cm})$  و  $(+3.0 \text{ cm}, +3.0 \text{ cm})$

احسب مقدار واتجاه محصلة المجال الكهربائي عند نقطة موقعها  $(0.0 \text{ cm}, +3.0 \text{ cm})$



According to the figure, if  $(Q = -2 \mu C)$  and  $(q = +2 \mu C)$ ,  
what is the electric field at the point P (0,0)


وفقاً للشكل، إذا  $(Q = -2 \mu C)$  و  $(q = +2 \mu C)$ ، ما المجال الكهربائي عند النقطة P (0,0)؟



According to the figure, two charges  $(Q = +8.0 \mu C)$  and  $(q = -5.0 \mu C)$ , the electric  
force between them equals to  $(0.36 N)$ , if Q is placed at  $(0.0 m, 0.8 m)$ .


What is the electric field at the point  $(0,0)$ ?

وفقاً للشكل، شحنتان  $(Q = +8.0 \mu C)$  و  $(q = -5.0 \mu C)$ ، القوة الكهربائية بينهما تساوي  $(0.36 N)$ ،  
إذا وُضعت Q عند النقطة  $(0.0 m, 0.8 m)$  ما هو المجال الكهربائي عند النقطة  $(0,0)$ ؟



Two identical point charges, each with a charge of  $+10 \mu\text{C}$ , are located along the  $x$ -axis at  $x = -2 \text{ cm}$  and  $x = 4 \text{ cm}$ , as shown below.

شحنتان نقطيتان متماثلتين، شحنة كل منهما  $+10 \mu\text{C}$ ، تقعان على طول المحور  $x$  عند  $x = -2 \text{ cm}$  و  $x = 4 \text{ cm}$ ، كما هو موضع أدناه.



What is the **magnitude and direction** of the net electric field at the origin (0 cm)?

ما مقدار واتجاه المجال الكهربائي الكلي عند نقطة الأصل (0 cm)؟

At what **location** along the  $x$ -axis will the electric field be equal to **zero**?

عند أي موضع على المحور  $x$  يساوي المجال الكهربائي صفر؟

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

Two charges,  $q_1 = -5.0 \mu\text{C}$  and  $q_2 = +4.0 \mu\text{C}$  are located in the  $xy$ -plane as shown in the figure below.

تقع الشحنتان  $q_1 = -5.0 \mu\text{C}$  و  $q_2 = +4.0 \mu\text{C}$  في المستوى  $xy$  كما هو موضع في الشكل التالي.

What is the **magnitude and direction** of the net electric field at origin?

ما مقدار واتجاه المجال الكهربائي الكلي عند نقطة الأصل (0 cm)؟



.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

Electric fields of varying magnitudes are directed either inward or outward at right angles on the faces of a cube, as shown in the figure. What is the strength and direction of the field on face  $F$ ?

توجه مجالات كهربائية بقادير مختلفه إلى الداخل أو الخارج بزاوية قائمة على الأوجه للمكعب، كما هو موضح في الشكل.

ما شدة واتجاه المجال على الوجه  $F$  ؟



.....

.....

.....

.....

.....

.....

.....

.....

According to the figure, a cube that has Volume  $0.001\text{m}^3$  in a uniform electric field ( $E=3 \times 10^{-2}\text{N/C}$ ), that is perpendicular to the plane of one face of the cube.

- What Is the magnitude of electric flux passing through the black face?
- What is the total electrical flux through that surface when charge  $q = 3.2\mu\text{C}$  place in the centre of cube

وفقاً للشكل، مكعب حجمه  $0.001\text{m}^3$  في مجال كهربائي منتظم ( $E=3 \times 10^{-2}\text{N/C}$ ) عمودي على مستوى أحد أوجه المكعب .

ما مقدار التدفق الكهربائي المار عبر الوجه الأسود؟

ما هو التدفق الكهربائي الكلي خلال هذا السطح عندما تكون الشحنة  $q = 3.2\mu\text{C}$  في مركز المكعب

.....

.....

.....


.....

.....

.....

.....

.....



The six faces of a cubical box each measure 20.0 cm by 20.0 cm, and the faces are numbered such that faces 1 and 6 are opposite to each other, as are faces 2 and 5, and faces 3 and 4. The flux through each face is given on the table. Find the net charge inside the cube

الأوجه الستة لصندوق أبعاد كل منها 20.0 في 20.0، والأوجه مرقمة؛ بحيث تكون الأوجه بحيث يكون الوجهان 1 و6 متقابلين، وكذلك الوجهان 2 و5، والوجهان 3 و4. التدفق خلال كل وجه معطى في الجدول. أوجد الشحنة الكلية داخل المكعب

| Face | Flux ( $\text{N m}^2/\text{C}$ ) |
|------|----------------------------------|
| 1    | -70.0                            |
| 2    | -300.0                           |
| 3    | -300.0                           |
| 4    | +300.0                           |
| 5    | -400.0                           |
| 6    | -500.0                           |

Prove that the electric flux through a closed surface is given by the net charge inside the surface divided by the permittivity of the medium, and write the Gauss's law in its integral form

برهن على أن التدفق الكهربائي خلال سطح مغلق يساوي الشحنة الكلية داخل السطح مقسومة على السماحية للوسط، واكتب قانون جاوس في صورته التكاملية

Find the net electric flux through the spherical closed surface shown below.

أوجد التدفق الكهربائي الكلي خلال السطح الكروي المغلق الموضح بالأسفل..

.....  
.....  
.....

+2.00 nC

+1.00 nC

-3.00 nC

According to the figure, a uniform electric field ( $E = 360\text{N/C}$ ), that is perpendicular to the plane of one face of the cube. if the electric flux passing through the left shaded face is equal to  $(-1.2\text{Nm}^2/\text{C})$ . What is the cube side **length**?

وفقاً للشكل، مجال كهربائي منتظم، ( $E = 360\text{N/C}$ ) عمودي على مستوى أحد أوجه المكعب، إذا كان التدفق الكهربائي المار بالوجه المظلل الأيسر يساوي  $(-1.2\text{Nm}^2/\text{C})$  ما طول ضلع المكعب؟

A point charge sits in the center of a cube. The electric flux through one side of the cube is  $7.2 \times 10^{10} \text{ Nm}^2/\text{C}$ . What is the charge at the center of the cube?

توجد شحنة نقطية في مركز مكعب. التدفق الكهربائي خلال أحد أضلاع المكعب يساوي  $7.2 \times 10^{10} \text{ Nm}^2/\text{C}$  ما الشحنة عند مركز المكعب؟

.....


.....

.....

.....

.....

.....



According to the figure, a uniform electric field ( $E = 28\text{N/C}$ ), that is perpendicular to the plane of one face of the cube. If the electric flux passing through the left shaded face is equal to  $(-7.0 \text{ Nm}^2/\text{C})$ , what is the volume the cube?

وفقاً للشكل، مجال كهربائي منتظم، ( $E = 28\text{N/C}$ ) عمودي على مستوى أحد أوجه المكعب. إذا كان التدفق الكهربائي المار بالوجه المظلل الأيسر يساوي  $(-7.0 \text{ Nm}^2/\text{C})$ ، فما حجم المكعب؟

A 3.5 C point charge sits in the center of a cube.

What is the electric flux through one side of the cube?

توجد شحنة نقطية 3.5 C في مركز مكعب. ما التدفق الكهربائي خلال أحد أضلاع المكعب؟

.....


.....

.....

.....

.....

.....





A positive charge of  $4.50\mu\text{C}$  is fixed in place. A particle of mass  $6.00\text{ g}$  and charge  $+3.00\mu\text{C}$  is fired with an initial speed of  $66.0\text{ m/s}$  directly toward the fixed charge from  $4.20\text{ cm}$  away. How close does the moving charge get to the fixed charge before it comes to rest and starts moving away from the fixed charge?

شحنة موجبة مقدارها  $4.50\mu\text{C}$  ثابتة في مكانها. وأطلق جسيم كتلته  $6.00\text{ g}$  وشحنته  $+3.00\mu\text{C}$  بسرعة ابتدائية مقدارها  $66.0\text{ m/s}$  مباشرة باتجاه الشحنة الثابتة من مسافة تبعد  $4.20\text{ cm}$  إلى أي مدى تقترب الشحنة المتحركة من الشحنة الثابتة قبل أن تصل إلى وضع السكون وتبدأ في الابتعاد عن الشحنة الثابتة؟


.....

.....

.....

.....

.....



A particle charge of  $(+2.0\mu\text{C})$  is released from rest at a point on the  $x$ -axis, ( $x = +6.0\text{ cm}$ ). It begins to move due to the presence of  $(+8.0\mu\text{C})$  a charge that remains fixed at the origin ( $x = 0.0$ ).

- Calculate the kinetic energy of the particle at the instant it passes the point ( $x = +15\text{ cm}$ ).

تحرر جسيم شحنته  $(+2.0\mu\text{C})$  من السكون عند نقطة على المحور  $x$  ( $x = +6.0\text{ cm}$ ). يبدأ في الحركة بسبب وجود شحنة  $(+8.0\mu\text{C})$  التي تظل ثابتة عند نقطة الأصل ( $x = 0.0$ ). احسب طاقة حركة الجسيم في اللحظة التي يمر فيها بالنقطة ( $x = +15\text{ cm}$ ).


.....

.....

.....

.....

.....



A negative electric charge  $-40.0\mu\text{C}$  is placed on the  $x$ -axis at point  $x = 2.0\text{ m}$ . Find the electric potential at  $x = 5.0\text{ m}$

الشحنة الكهربائية السالبة  $-40.0\mu\text{C}$  على المحور  $x$  عند النقطة  $x = 2.0\text{ m}$ . أوجد الجهد الكهربائي عند  $x = 5.0\text{ m}$


.....

.....

.....

.....

.....



As shown in Figure a charge  $q_1 = 2.00 \mu\text{C}$  is located at the origin and a charge  $q_2 = 26.00 \mu\text{C}$  is located at  $(0, 3.00) \text{ m}$ .


Find the total electric potential due to these charges at point  $P$ , whose coordinates are  $(4.00, 0) \text{ m}$ .

كما هو موضح في الشكل الشحنة  $q_1 = 2.00 \mu\text{C}$  تقع عند نقطة الأصل وشحنة  $q_2 = 26.00 \mu\text{C}$  عند النقطة  $(0, 3.00) \text{ m}$ . أوجد الجهد الكهربائي الكلي الناتج عن هذه الشحنات عند النقطة  $P$ ، التي إحداثياتها  $(4.00, 0) \text{ m}$ .

$$V_P = k_e \left( \frac{q_1}{r_1} + \frac{q_2}{r_2} \right)$$

$$V_P = (9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2) \left( \frac{2.00 \times 10^{-6} \text{ C}}{4.00 \text{ m}} + \frac{-6.00 \times 10^{-6} \text{ C}}{5.00 \text{ m}} \right)$$

$$= -6.29 \times 10^3 \text{ V}$$




Find the change in potential energy of the system of two charges plus a third charge  $q_3 = 3.00 \mu\text{C}$  as the charge moves from infinity to point  $P$ .

أوجد التغير في طاقة وضع النظام المكون من شحنتين شحنة ثالثة  $q_3 = 3.0 \mu\text{C}$  عندما تتحرك الشحنة من اللانهاية إلى

$$U_f = q_3 V_P$$

$$\Delta U = U_f - U_i = q_3 V_P - 0 = (3.00 \times 10^{-6} \text{ C})(-6.29 \times 10^3 \text{ V})$$

$$= -1.89 \times 10^{-2} \text{ J}$$



system of point charges. shows three-point charges

$q_1 = +1.50 \mu\text{C}$ ,  $q_2 = +2.50 \mu\text{C}$ , and  $q_3 = -3.50 \mu\text{C}$ .

Charge  $q_1$  is located at  $(0, a)$ ,  $q_2$  is located at  $(0,0)$ , and  $q_3$  is located at  $(b,0)$ , where  $a =$

$8.00 \text{ m}$  and  $b = 6.00 \text{ m}$ . The electric potential at point  $P$


يوضِّح الشحنات النقطية الثلاثي  $q_1 = +1.50 \mu\text{C}$ ,  $q_2 = +2.50 \mu\text{C}$ , and  $q_3 = -3.50 \mu\text{C}$ .

تقع الشحنة  $q_1$  عند  $(0, a)$ ، وتقع الشحنة  $q_2$  عند  $(0, 0)$ ، وتقع الشحنة  $q_3$  عند  $(b, 0)$  عند  $a = 8.00 \text{ m}$  و  $b = 6.00 \text{ m}$ . الجهد الكهربائي عند النقطة

$$V = \sum_{i=1}^3 \frac{kq_i}{r_i} = k \left( \frac{q_1}{r_1} + \frac{q_2}{r_2} + \frac{q_3}{r_3} \right) = k \left( \frac{q_1}{b} + \frac{q_2}{\sqrt{a^2 + b^2}} + \frac{q_3}{a} \right)$$


$$= (8.99 \times 10^9 \text{ N m}^2/\text{C}^2) \left( \frac{1.50 \times 10^{-6} \text{ C}}{6.00 \text{ m}} + \frac{2.50 \times 10^{-6} \text{ C}}{\sqrt{(8.00 \text{ m})^2 + (6.00 \text{ m})^2}} + \frac{-3.50 \times 10^{-6} \text{ C}}{8.00 \text{ m}} \right)$$

$$= 562 \text{ V}$$



two-point charges are located at two corners of a rectangle, as shown in the figure

- What is the electric potential at point A?
- What is the potential difference between points A and B?  
توجد شحنتان عند زاويتين في مستطيل، كما هو موضح في الشكل  
ما الجهد الكهربائي عند النقطة (A)؟ ما فرق الجهد بين النقطتين (A) و (B)؟



$$V_A = \frac{kq_1}{r_1} + \frac{kq_2}{r_2} = k \left( \frac{q_1}{r_1} + \frac{q_2}{r_2} \right)$$

$$V_A = (9.0 \times 10^9 \text{ Nm}^2/\text{C}^2) \left( \frac{-1.00 \times 10^{-6} \text{ C}}{0.250 \text{ m}} + \frac{3.00 \times 10^{-6} \text{ C}}{0.500 \text{ m}} \right) = 1.8 \times 10^4 \text{ V}$$

$$V_{AB} = V_A - V_B = k \left( \frac{q_1}{r_1} + \frac{q_2}{r_2} \right) - k \left( \frac{q_1}{r_2} + \frac{q_2}{r_1} \right) = k(q_1 - q_2) \left( \frac{1}{r_1} - \frac{1}{r_2} \right)$$

$$V_{AB} = (9.0 \times 10^9) (-1.00 \times 10^{-6} \text{ C} - 3.00 \times 10^{-6} \text{ C}) \left( \frac{1}{0.250} - \frac{1}{0.500} \right)$$


$$V_{AB} = 7.2 \times 10^4 \text{ V}$$

In the figure shown, charges  $q_1 = +2.0 \mu\text{C}$ ,  $q_2 = +4.0 \mu\text{C}$ , and  $Q = -3.0 \mu\text{C}$  are placed next to each other. The charges in an air.

Calculate the change in the electric potential energy of a charge when it is moved from point a to point b.


$q_1 = +2.0 \mu\text{C}$ ,  $q_2 = +4.0 \mu\text{C}$ ,  $Q = -3.0 \mu\text{C}$  في الشكل المجاور وضعت الشحنتان  $q_1 = +2.0 \mu\text{C}$ ،  $q_2 = +4.0 \mu\text{C}$ ، و  $Q = -3.0 \mu\text{C}$

وكان الهواء يحيط بالشحنتين. احسب التغير في طاقة الوضع الكهربائية لشحنة عندما تنقل من النقطة (a) إلى النقطة (b)



The point charge ( $q$ ) is placed in the air as in the adjacent figure. If the magnitude of the electric field strength at the point is ( $400V/m$ ) and the voltage at the point is ( $500V$ ), calculate the magnitude of the charge ( $q$ )

وضعت الشحنة النقطية ( $q$ ) في الهواء كما في الشكل المجاور ، فإذا كان مقدار شدة المجال الكهربائي عند النقطة ( $400V/m$ ) والجهد الكهربائي عندها ( $500V$ ) اصعب مقدار الشحنة ( $q$ )



.....

.....

.....

.....

.....

.....

.....

.....

As shown in the figure, if the magnitude of the electric potential at point O due to a point electric charge  $q$  is equal to  $+9 \times 10^{-2}V$

وفقا للشكل المقابل، إذا كان مقدار الجهد الكهربائي في النقطة O الناشي عن شحنة كهربائية نقطية  $q$  يساوي  $+9 \times 10^{-2}V$

A. Calculate the magnitude of the **charge**

اصعب مقدار الشحنة

B. The magnitude of **the electric field** at point O?

مقدار المجال الكهربائي في النقطة O؟

C. If you place a second point charge at point O, does the magnitude of the electric field at point O change

**Explain your answer.**

في حال وضعت شحنة نقطية أخرى في النقطة O، هل يتغير مقدار المجال الكهربائي في النقطة O **فسر اجابتك.**



.....

.....

.....

.....

.....

.....

.....

.....

This revision is designed to help you reinforce your understanding and grasp of the concepts we've covered throughout the term. Please remember that revising thoroughly requires going over all the materials presented in this term. Focusing on the core topics, examples, and exercises we discussed will strengthen your foundation and help you achieve the best possible results.

تم تصميم هذه المراجعة لمساعدتكم على تعزيز فهمكم وتثبيت المعلومات التي تناولناها خلال هذا الفصل. تذكروا أن المراجعة الفعالة تتطلب دراسة جميع الدروس التي تم تقديمها خلال هذا الفصل الدراسي. التركيز على المواضيع الأساسية، والأمثلة، والتمارين التي ناقشناها سيعزز أساسياتكم ويؤهلكم للحصول على أفضل النتائج الممكنة.

استاذ عبدالرحمن عصام عبيد