

شكراً لتحميلك هذا الملف من موقع المناهج الإماراتية



شرح وقوانين مفيدة (مناهج إنجليزية)

[موقع المناهج](#) ⇨ [المناهج الإماراتية](#) ⇨ [الصف التاسع المتقدم](#) ⇨ [رياضيات](#) ⇨ [الفصل الثالث](#) ⇨ [الملف](#)

تاريخ نشر الملف على موقع المناهج: 20:21:37 2019-06-16

التواصل الاجتماعي بحسب الصف التاسع المتقدم



روابط مواد الصف التاسع المتقدم على تلغرام

[الرياضيات](#)

[اللغة الانجليزية](#)

[اللغة العربية](#)

[التربية الاسلامية](#)

المزيد من الملفات بحسب الصف التاسع المتقدم والمادة رياضيات في الفصل الثالث

[حل أسئلة الامتحان النهائي الالكتروني ريفيل](#)

1

[أسئلة الامتحان النهائي الالكتروني بريدج](#)

2

[أسئلة الامتحان النهائي الورقي بريدج](#)

3

[حل أسئلة الاختبار التحريبي ريفيل](#)

4

[أسئلة نموذج تدريبي ريفيل](#)

5

كل ما يحتاجه الطالب في جميع الصفوف من أوراق عمل واختبارات ومذكرات، يجده هنا في الروابط التالية لأفضل مواقع تعليمي إماراتي 100 %

<u>تطبيق المناهج الإماراتية</u>	<u>الاجتماعيات</u>	<u>الرياضيات</u>
<u>الصفحة الرسمية على التلغرام</u>	<u>الاسلامية</u>	<u>العلوم</u>
<u>الصفحة الرسمية على الفيسبوك</u>	<u>الانجليزية</u>	
<u>التربية الاخلاقية لجميع الصفوف</u>	<u>اللغة العربية</u>	
<u>التربية الرياضية</u>		
مجموعات التلغرام.	مجموعات الفيسبوك	قنوات تلغرام
<u>الصف الأول</u>	<u>الصف الأول</u>	<u>الصف الأول</u>
<u>الصف الثاني</u>	<u>الصف الثاني</u>	<u>الصف الثاني</u>
<u>الصف الثالث</u>	<u>الصف الثالث</u>	<u>الصف الثالث</u>
<u>الصف الرابع</u>	<u>الصف الرابع</u>	<u>الصف الرابع</u>
<u>الصف الخامس</u>	<u>الصف الخامس</u>	<u>الصف الخامس</u>
<u>الصف السادس</u>	<u>الصف السادس</u>	<u>الصف السادس</u>
<u>الصف السابع</u>	<u>الصف السابع</u>	<u>الصف السابع</u>
<u>الصف الثامن</u>	<u>الصف الثامن</u>	<u>الصف الثامن</u>
<u>الصف التاسع عام</u>	<u>الصف التاسع عام</u>	<u>الصف التاسع عام</u>
<u>الصف التاسع متقدم</u>	<u>الصف التاسع متقدم</u>	<u>الصف التاسع متقدم</u>
<u>الصف العاشر عام</u>	<u>الصف العاشر عام</u>	<u>الصف العاشر عام</u>
<u>الصف العاشر متقدم</u>	<u>الصف العاشر متقدم</u>	<u>الصف العاشر متقدم</u>
<u>الحادي عشر عام</u>	<u>الحادي عشر عام</u>	<u>الحادي عشر عام</u>
<u>الحادي عشر متقدم</u>	<u>الحادي عشر متقدم</u>	<u>الحادي عشر متقدم</u>
<u>ثاني عشر عام</u>	<u>الثاني عشر عام</u>	<u>الثاني عشر عام</u>
<u>ثاني عشر متقدم</u>	<u>الثاني عشر متقدم</u>	<u>الثاني عشر متقدم</u>



Grade 9 Advanced Mathematics – Term 2 Revision Notes


Square of a Sum	$(a + b)^2 = a^2 + 2ab + b^2$ (a.k.a. as a perfect square)	
Square of a Difference	$(a - b)^2 = a^2 - 2ab + b^2$ (a.k.a. as a perfect square)	
Product of a Sum and a Difference	$(a + b)(a - b) = a^2 - b^2$ (a.k.a. as a <u>difference</u> of two squares)	<p style="text-align: center;"><u>Very useful for Factoring</u></p> $x^2 - 9 = (x + 3)(x - 3)$ $9n^2 - 4 = (3n + 2)(3n - 2)$ $16c^2 - 49d^2 = (4c + 7d)(4c - 7d)$ $x^4 - 25 = (x^2 + 5)(x^2 - 5)$
Test for Perfect Square and Factoring $a^2 + 2ab + b^2 = (a + b)^2$ $a^2 - 2ab + b^2 = (a - b)^2$ <ul style="list-style-type: none"> The first and last terms must be perfect squares The middle term must be twice the product of the square roots of the first and last terms 		
$16x^2 + 24x + 9$ <ul style="list-style-type: none"> $\sqrt{16x^2} = 4x$ and $\sqrt{9} = 3$ (both perfect squares) $2(4x)(3) = 24x$ (this is the middle term) So this is a perfect square We can factorise $16x^2 + 24x + 9 = (4x + 3)^2$ 	$9x^2 - 12x + 4$ <ul style="list-style-type: none"> $\sqrt{9x^2} = 3x$ and $\sqrt{4} = 2$ (both perfect squares) $-2(3x)(2) = -12x$ (this is the middle term) So this is a perfect square We can factorise $9x^2 - 12x + 4 = (3x - 2)^2$ 	$25x^2 + 20x + 9$ <ul style="list-style-type: none"> $\sqrt{25x^2} = 5x$ and $\sqrt{9} = 3$ (both perfect squares) $2(5x)(3) = 30x$ this is NOT the middle term not a perfect square <hr/> $4a^2 - 4a + 2$ <ul style="list-style-type: none"> $\sqrt{2}$ is not a perfect square So $4a^2 - 4a + 2$ is not a perfect square
Factoring Polynomials by Grouping	Polynomials with four or more terms. $ax + bx + ay + by$ $= x(a + b) + y(a + b)$ $= (a + b)(x + y)$	$4qr + 8r + 3q + 6$ $= (4qr + 8r) + (3q + 6)$ <div style="display: flex; justify-content: space-between; font-size: small;"> group terms with common factors factor the GCF from each group </div> $= 4r(q + 2) + 3(q + 2)$ <div style="display: flex; justify-content: space-between; font-size: small;"> factor the GCF from each group distributive property </div> $= (q + 2)(4r + 3)$ <hr/> $2mk - 12m + 42 - 7k$ $= (2mk - 12m) + (42 - 7k)$ <div style="display: flex; justify-content: space-between; font-size: small;"> group terms with common factors factor the GCF from each group </div> $= 2m(k - 6) - 7(k - 6)$ <div style="display: flex; justify-content: space-between; font-size: small;"> factor the GCF from each group distributive property </div> $= (k - 6)(2m - 7)$

Factoring Trinomial Quadratic Expressions



(the more practise the easier this will become)

$x^2 + bx + c$	$ax^2 + bx + c$																				
<p>Find m and p with a product of c and a sum of b. i.e. $m \cdot p = c$ and $m + p = b$ then re-write $x^2 + bx + c$ as $(x + m)(x + p)$</p>	<p>Find m and p with a product of ac and a sum of b. i.e. $m \cdot p = ac$ and $m + p = b$ then re-write $ax^2 + bx + c$ as $ax^2 + mx + px + c$ finally factor by grouping.</p>																				
<p>$x^2 + 7x + 12$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of 12</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>1, 12</td> <td>13</td> </tr> <tr> <td>2, 6</td> <td>8</td> </tr> <tr> <td>3, 4</td> <td>7</td> </tr> </tbody> </table> <p>b and c are both +ve so factors of 12 will be +ve</p> <p>$(x + 3)(x + 4)$</p>	Factors of 12	Sum of Factors	1, 12	13	2, 6	8	3, 4	7	<p>$2x^2 + 5x + 3$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of ac</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>1, 6</td> <td>7</td> </tr> <tr> <td>2, 3</td> <td>5</td> </tr> </tbody> </table> <p>$ac = 6$</p> <p>$2x^2 + 2x + 3x + 3$</p> <p>$2x(x + 1) + 3(x + 1)$</p> <p>$(x + 1)(2x + 3)$</p>	Factors of ac	Sum of Factors	1, 6	7	2, 3	5						
Factors of 12	Sum of Factors																				
1, 12	13																				
2, 6	8																				
3, 4	7																				
Factors of ac	Sum of Factors																				
1, 6	7																				
2, 3	5																				
<p>$x^2 - 22x + 21$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of 21</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>-1, -21</td> <td>-22</td> </tr> <tr> <td>-3, -7</td> <td>-10</td> </tr> </tbody> </table> <p>b is -ve and c is +ve so factors of 21 will be -ve</p> <p>$(x - 1)(x - 21)$</p>	Factors of 21	Sum of Factors	-1, -21	-22	-3, -7	-10	<p>$3x^2 - 17x + 20$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of ac</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>-1, -60</td> <td>-61</td> </tr> <tr> <td>-2, -30</td> <td>-32</td> </tr> <tr> <td>-3, -20</td> <td>-23</td> </tr> <tr> <td>-4, -15</td> <td>-19</td> </tr> <tr> <td>-5, -12</td> <td>-17</td> </tr> <tr> <td>-6, -10</td> <td>-16</td> </tr> </tbody> </table> <p>$ac = 60$</p> <p>$3x^2 - 5x - 12x + 20$</p> <p>$x(3x - 5) - 4(3x - 5)$</p> <p>$(3x - 5)(x - 4)$</p>	Factors of ac	Sum of Factors	-1, -60	-61	-2, -30	-32	-3, -20	-23	-4, -15	-19	-5, -12	-17	-6, -10	-16
Factors of 21	Sum of Factors																				
-1, -21	-22																				
-3, -7	-10																				
Factors of ac	Sum of Factors																				
-1, -60	-61																				
-2, -30	-32																				
-3, -20	-23																				
-4, -15	-19																				
-5, -12	-17																				
-6, -10	-16																				
<p>$x^2 + 2x - 15$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of -15</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>-1, +15</td> <td>14</td> </tr> <tr> <td>-3, +5</td> <td>2</td> </tr> </tbody> </table> <p>c is -ve so factors of -15 have opposite signs (since b is +ve the greater absolute value factor is also +ve)</p> <p>$(x - 3)(x + 5)$</p>	Factors of -15	Sum of Factors	-1, +15	14	-3, +5	2	<p>$2x^2 + 3x - 5$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of ac</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>-1, +10</td> <td>9</td> </tr> <tr> <td>-2, +5</td> <td>3</td> </tr> </tbody> </table> <p>$ac = -10$</p> <p>$2x^2 - 2x + 5x - 5$</p> <p>$2x(x - 1) + 5(x - 1)$</p> <p>$(x - 1)(2x + 5)$</p>	Factors of ac	Sum of Factors	-1, +10	9	-2, +5	3								
Factors of -15	Sum of Factors																				
-1, +15	14																				
-3, +5	2																				
Factors of ac	Sum of Factors																				
-1, +10	9																				
-2, +5	3																				
<p>$x^2 - 7x - 18$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of -18</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>+1, -18</td> <td>-17</td> </tr> <tr> <td>+2, -9</td> <td>-7</td> </tr> <tr> <td>+3, -6</td> <td>-3</td> </tr> </tbody> </table> <p>c is -ve so factors of -18 have opposite signs (since b is -ve the greater absolute value factor is also -ve)</p> <p>$(x + 2)(x - 9)$</p>	Factors of -18	Sum of Factors	+1, -18	-17	+2, -9	-7	+3, -6	-3	<p>$6x^2 - 7x - 3$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of ac</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>+1, -18</td> <td>-17</td> </tr> <tr> <td>+2, -9</td> <td>-7</td> </tr> <tr> <td>+3, -6</td> <td>-3</td> </tr> </tbody> </table> <p>$ac = -18$</p> <p>$6x^2 + 2x - 9x - 3$</p> <p>$2x(3x + 1) - 3(3x + 1)$</p> <p>$(3x + 1)(2x - 3)$</p>	Factors of ac	Sum of Factors	+1, -18	-17	+2, -9	-7	+3, -6	-3				
Factors of -18	Sum of Factors																				
+1, -18	-17																				
+2, -9	-7																				
+3, -6	-3																				
Factors of ac	Sum of Factors																				
+1, -18	-17																				
+2, -9	-7																				
+3, -6	-3																				
Determine if a Polynomial is Prime	Zero Product Property																				
<p>$4x^2 - 3x + 5$</p> <p>$ac = 20$</p> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left; border-bottom: 1px solid black;">Factors of ac</th> <th style="text-align: left; border-bottom: 1px solid black;">Sum of Factors</th> </tr> </thead> <tbody> <tr> <td>-1, -20</td> <td>-21</td> </tr> <tr> <td>-2, -10</td> <td>-12</td> </tr> <tr> <td>-4, -5</td> <td>-9</td> </tr> </tbody> </table> <p>There are <u>no factors</u> with a sum of -3. So the quadratic expression cannot be factored using integers. Therefore this is prime.</p>	Factors of ac	Sum of Factors	-1, -20	-21	-2, -10	-12	-4, -5	-9	<p>If the produce of two factors is 0, then at least one of the factors must be 0. If $ab = 0$, then $a = 0$ or $b = 0$ or both a and b equal 0.</p> <p>$(2d + 6)(3d - 15) = 0$</p> <p>$2d + 6 = 0$ or $3d - 15 = 0$</p> <p>$2d = -6$ $3d = 15$</p> <p>$d = -3$ $d = 5$</p>												
Factors of ac	Sum of Factors																				
-1, -20	-21																				
-2, -10	-12																				
-4, -5	-9																				

Graphing Exponential Growth	$y = ab^x$ $a > 0$ $b > 1$ <p>(<i>a</i> is the <i>y</i> – intercept)</p>	
Graphing Exponential Decay	$y = ab^x$ $a > 0$ $0 < b < 1$ <p>(<i>a</i> is the <i>y</i> – intercept)</p>	
Exponential Growth	$y = a(1 + r)^t$ <p> <i>y</i> = final amount <i>a</i> = initial amount <i>r</i> = rate of change (decimal) <i>t</i> = time </p>	
Exponential Decay	<p style="text-align: center;">http://alManahj.com/ae</p> $y = a(1 - r)^t$ <p> <i>y</i> = final amount <i>a</i> = initial amount <i>r</i> = rate of change (decimal) <i>t</i> = time </p>	
Compound Interest	$A = P \left(1 + \frac{r}{n} \right)^{nt}$ <p> <i>A</i> = current amount <i>P</i> = principal amount (initial) <i>r</i> = annual interest rate (decimal) <i>n</i> = no. of times interest is compounded <i>t</i> = time in years </p>	<p> compounded monthly <i>n</i> = 12 compounded quarterly <i>n</i> = 4 compounded daily <i>n</i> = 365 </p>
Geometric Sequence	<p style="text-align: center;"><i>n</i>th term</p> $a_n = a_1 r^{n-1}$ <p> <i>a</i>₁ = first term <i>r</i> = common ratio (2nd term divide 1st term) </p>	<p style="text-align: center;">Recursive Formula</p> $a_n = r \cdot a_{n-1}$ <p>(Remember <i>a</i>_{<i>n</i>-1} is the previous term)</p>

<p>Arithmetic Sequence</p>	<p>n^{th} term</p> $a_n = a_1 + (n - 1)d$ <p>$a_1 =$ first term $d =$ common difference</p>	<p>Recursive Formula</p> $a_n = a_{n-1} + d$ <p>(Remember a_{n-1} is the previous term)</p>
<p>Graphing Radical Function</p>	$y = \sqrt{x}$ <p>(remember the radicand can't be negative)</p>	
<p>Transformation of Radical Function 1 Dilation</p>	$y = a\sqrt{x}$ $a > 1$ <p>Parent function: $y = \sqrt{x}$</p> <p>(stretched vertically)</p>	$y = a\sqrt{x}$ $0 < a < 1$ <p>Parent function: $y = \sqrt{x}$</p> <p>(compressed vertically)</p>

<http://alManahj.com/ae>




Transformation of
Radical Function 2
Reflection

$$y = a\sqrt{x}$$

$$a < 0$$

$$\text{Parent function: } y = \sqrt{x}$$

(reflection across x - axis)



Transformation of
Radical Function 3
Reflection

$$y = \sqrt{x + h} + k$$

Translate the graph k units UP if $k > 0$ and $|k|$ units DOWN if $k < 0$

Translate the graph h units LEFT if $h > 0$ and $|h|$ units RIGHT if $h < 0$

$$\text{Parent function: } y = \sqrt{x}$$

Simplifying a Square Root with Variables


<p>Variables with EVEN exponents in radicand and the simplified answer has an EVEN exponent</p> $\sqrt{x^4} = x^2$ $\sqrt{x^8} = x^4$ $\sqrt{x^{12}} = x^6$	<p>Variables with EVEN exponents in radicand and the simplified answer has an ODD exponent (must use ABSOLUTE VALUE symbol)</p> $\sqrt{x^2} = x $ $\sqrt{x^6} = x^3 $ $\sqrt{x^{22}} = x^{11} $	<p>Variables with ODD exponents in radicand</p> $\sqrt{x^5} = \sqrt{x^4} \sqrt{x} = x^2 \sqrt{x}$ $\sqrt{x^{13}} = \sqrt{x^{12}} \sqrt{x} = x^6 \sqrt{x}$ $\sqrt{x^{15}} = \sqrt{x^{14}} \sqrt{x} = x^7 \sqrt{x}$
--	---	--


Direct Variation	$y = kx$	k is known as the <i>constant of variation</i> or <i>constant of proportionality</i>
-------------------------	----------	--


Inverse Variation	$xy = k$ $x \neq 0$ and $y \neq 0$	k is a non-zero constant
--------------------------	---------------------------------------	-----------------------------------


Product Rule for Inverse Variation	$x_1 y_1 = x_2 y_2$	If (x_1, y_1) and (x_2, y_2) are solutions of an inverse variation then the products $x_1 y_1$ and $x_2 y_2$ are equal.
---	---------------------	--





Excluded Values for a Rational Function ($y = \frac{1}{x}$)	Division by zero is undefined . Any value that results in a denominator of zero is excluded from the domain of a rational function.	$y = \frac{5}{4x - 8}$ $4x - 8 = 0$ $4x = 8$ $x = 2$ <p>The excluded value is $x = 2$</p>
--	--	--

Asymptotes	$y = \frac{a}{x - b} + c$ $a \neq 0$ <p>vertical asymptote: $x = b$ horizontal asymptote: $y = c$</p>	
-------------------	--	---

Point	A point is a location. It has neither shape nor size. point A	
--------------	--	---

Line	There is exactly one line through any two points . line m, line PQ or \overleftrightarrow{PQ}, line QP or \overleftrightarrow{QP}	
-------------	---	---

Plane	There is exactly one plane through any three points that are not collinear . plane \mathcal{K}, plane BCD, plane BDC, plane CBD, plane CDB, plane DBC, plane DCB	
--------------	---	---

Collinear	Points that lie on the same line .	
Coplanar	Points that lie in the same plane .	
Segment	<p>A line segment can be measured because it has two end points.</p> <p>Label: \overline{AB} or \overline{BA} Measure: AB or BA</p>	
Congruent segments	<p>Congruent segments have the same measure.</p> <p>\cong "is congruent to"</p> <p>$\overline{AB} \cong \overline{CD}$ which means $AB = CD$</p>	
Distance Formula	<p>If P has coordinates (x_1, y_1) and Q has coordinates (x_2, y_2) then</p> $PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$	
Midpoint Formula	<p>If \overline{PQ} has midpoints $P(x_1, y_1)$ and $Q(x_2, y_2)$ in the coordinate plane, then the midpoint M of \overline{PQ} has coordinates</p> $M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$	

<http://alManahj.com/ae>

Truth Tables

Negation	
p	$\sim p$
T	F
F	T

"and"		
Conjunction		
p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

"or"		
Disjunction		
p	q	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Conditional Statement	$p \rightarrow q$ read as if <i>p then q</i> or <i>p implies q</i> p is the hypothesis q is the conclusion	<table border="1"> <thead> <tr> <th colspan="3">Conditional Statements</th> </tr> <tr> <th>p</th> <th>q</th> <th>$p \rightarrow q$</th> </tr> </thead> <tbody> <tr> <td>T</td> <td>T</td> <td>T</td> </tr> <tr> <td>T</td> <td>F</td> <td>F</td> </tr> <tr> <td>F</td> <td>T</td> <td>T</td> </tr> <tr> <td>F</td> <td>F</td> <td>T</td> </tr> </tbody> </table>	Conditional Statements			p	q	$p \rightarrow q$	T	T	T	T	F	F	F	T	T	F	F	T
	Conditional Statements																			
p	q	$p \rightarrow q$																		
T	T	T																		
T	F	F																		
F	T	T																		
F	F	T																		
Related Conditionals																				
$p \rightarrow q$	Conditional	Logically Equivalent Statements <ul style="list-style-type: none"> A <i>conditional</i> and its <i>contrapositive</i> are logically equivalent The <i>converse</i> and <i>inverse</i> of a conditional are logically equivalent 																		
$q \rightarrow p$	Converse																			
$\sim p \rightarrow \sim q$	Inverse																			
$\sim q \rightarrow \sim p$	Contrapositive																			
Inductive Reasoning	Inductive reasoning uses patterns and observations to make a conjecture.																			
Deductive Reasoning	Deductive reasoning uses facts, rules, definitions and properties to make a conjecture.																			
Law of Detachment	If $p \rightarrow q$ and p is true, then q is true.	Helps us determine if a conclusion is valid or not.																		
Law of Syllogism	If $p \rightarrow q$ and $q \rightarrow r$ are true then $p \rightarrow r$ is true.																			

<http://alManahj.com/ae>